arXiv:math/0612216v2 [math.CA] 28 Mar 2007

# PLANCHEREL-ROTACH ASYMPTOTICS FOR $q$-SERIES

RUIMING ZHANG

Abstract. In this work we study the Plancherel-Rotach type asymptotics for selected $q$-series and $q$-orthogonal polynomials with complex scalings. The $q$-series we cover are Euler's $q$-exponential, Ramanujan function, Jackson's $q$-Bessel function of second kind, Ismail-Masson orthogonal polynomials, Stieltjes-Wigert polynomials and $q$-Laguerre polynomials. For a fixed $q$ with $0 < q < 1$, in each case the main term of the asymptotic formulas may contain Ramanujan function or theta function depending on the value of scaling parameter. Furthemore, when the scaling parameter is in certain strip of the complex plane, its number theoretical property completely determines the order of the error term. In each cases, we also investigate the asymptotic behavior of the mentioned $q$-series when $q$ approaching 1 in a restricted manner. These asymptotic formulas may provide insights to new random matrix models.

## 1. Introduction

The Plancherel-Rotach asymptotics for classical orthogonal polynomials are essential to obtain universality results in random matrix theory [21]. The associated random matrix models for $q$-orthogonal polynomials are still unknown today. It might be interesting to calculate the Plancherel-Rotach type asymptotics for $q$-orthogonal polynomials to gain some insights to the related random matrix models.

The Plancherel-Rotach type asymptotics for $q$-orthogonal polynomials started in [18], where we studied Plancherel-Rotach type asymptotics for three families of $q$-orthogonal polynomials with real logarithmic scalings. They are Ismail-Masson polynomials $\{h_n(x|q)\}_{n=0}^{\infty}$, Stieltjes-Wigert polynomials $\{S_n(x;q)\}_{n=0}^{\infty}$ and $q$-Laguerre polynomials $\left\{L_n^{(\alpha)}(x;q)\right\}_{n=0}^{\infty}$. These $q$-orthogonal polynomials are associated with indeterminant moment problems [1]. The asymptotics reveal a remarkable pattern which is quite different to the pattern associated with classical Plancherel-Rotach asymptotics [25, 13, 7, 8]. The main term of asymptotics may contain Ramanujan function $A_q(z)$ or theta function according to the value of the scaling parameter.

In this work we will investigate Plancherel-Rotach type asymptotics with complex scalings for some $q$-series, first with fixed $q \in (0,1)$, then let $q$ approaching 1 in some restricted manners. The $q$-series treated in this work are Euler's $q$-exponential $E_q(z)$, Ramanujan function $A_q(z)$, Jackson's $q$-Bessel functions of second kind $J_\nu^{(2)}(z;q)$, Ismail-Masson orthogonal polynomials $\{h_n(x|q)\}_{n=0}^{\infty}$, Stieltjes-Wigert polynomials $\{S_n(x;q)\}_{n=0}^{\infty}$ and $q$-Laguerre polynomials $\left\{L_n^{(\alpha)}(x;q)\right\}_{n=0}^{\infty}$.

---

1991 *Mathematics Subject Classification.* Primary 30E15. Secondary 33D45.

*Key words and phrases.* $q$-Orthogonal polynomials, Euler's $q$-Exponential, Ramanujan function, $q$-Gamma function, Jackson's $q$-Bessel function Ismail-Masson orthogonal polynomials, Stieltjes-Wigert orthogonal polynomials, $q$-Laguerre orthogonal polynomials, Plancherel-Rotach asymptotics, theta functions, complex scaling, random matrix, evaluation of definite integrals.

The results reveal an even more complicated pattern. The error terms manifest a link between special functions and number theory. The method we use in this study is a discrete version of the classical Laplace method in asymptotics [18]. This version of discrete Laplace method may also be applied to study the asymptotics for other $q$-series such as general confluent basic hypergeometric functions. We prefer to publish these results at another time.

Through out this work, We shall assume that $0 < q < 1$ unless otherwise stated. We also assume that $s = \sigma + it$ is a complex number with $\sigma = \Re(s)$ and $t = \Im(s)$. All the log and power functions are taken as their principle branches. The asymptotic formulas here are mostly for conceptual purposes, so we won't pursue the best orders nor the best implied constants.

1.1. **$q$-series.** For any complex number $a$, we define [6, 9, 13, 20]

$$(a;q)_\infty := \prod_{k=0}^{\infty}(1-aq^k) \tag{1}$$

and the $q$-shifted factorial of $a, a_1, \dots a_m$ are defined by

$$(a;q)_n := \frac{(a;q)_\infty}{(aq^n;q)_\infty}, \quad (a_1,\dots,a_m;q)_n := \prod_{k=1}^{m}(a_k;q)_n \tag{2}$$

for all integers $n \in \mathbb{Z}$ amd $m \in \mathbb{N}$.

**Lemma 1.1.** *Given any complex number $z$, we assume that*

$$0 < \frac{|z|\,q^n}{1-q} < \frac{1}{2} \tag{3}$$

*for some positive integer $n$. Then,*

$$\frac{(z;q)_\infty}{(z;q)_n} = (zq^n;q)_\infty := 1 + r_1(z;n), \tag{4}$$

*with*

$$|r_1(z;n)| \le \frac{2\,|z|\,q^n}{1-q}, \tag{5}$$

*and*

$$\frac{(z;q)_n}{(z;q)_\infty} = \frac{1}{(zq^n;q)_\infty} := 1 + r_2(z;n), \tag{6}$$

*with*

$$|r_2(z;n)| \le \frac{2\,|z|\,q^n}{(1-q)}. \tag{7}$$

*Proof.* From the $q$-binomial theorem [6, 9, 13, 20]

$$\frac{(Az;q)_\infty}{(z;q)_\infty} = \sum_{k=0}^{\infty}\frac{(A;q)_k}{(q;q)_k}z^k, \quad A, z \in \mathbb{C}, \quad |z| < 1, \tag{8}$$

and the inequality

$$(q;q)_k \ge (1-q)^k \tag{9}$$

for $k = 0, 1, ...$ we obtain

$$r_2(z;n) \quad = \quad \sum_{k=0}^{\infty} \frac{(zq^n)^{k+1}}{(q;q)_{k+1}}, \tag{10}$$

and

$$|r_2(z;n)| \le \sum_{k=0}^{\infty} \frac{(|z|\, q^n)^{k+1}}{(q;q)_{k+1}} \le \frac{|z|\, q^n}{(1-q)} \sum_{k=0}^{\infty} \left(\frac{|z|\, q^n}{1-q}\right)^k \le \frac{2\,|z|\, q^n}{(1-q)}. \tag{11}$$

Apply a limiting case of (8),

$$(z;q)_\infty = \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{(q;q)_k}(-z)^k \quad z \in \mathbb{C}, \tag{12}$$

and the inequalities,

$$\frac{1-q^k}{1-q} \ge kq^{k-1}, \quad \frac{(q;q)_k}{(1-q)^k} \ge k!q^{k(k-1)/2} \tag{13}$$

for $k = 0, 1, \dots$ we get

$$r_1(z;n) = \sum_{k=1}^{\infty} \frac{q^{k(k-1)/2}(-zq^n)^k}{(q;q)_k}, \tag{14}$$

and

$$\begin{aligned} |r_1(z;n)| &\le \sum_{k=0}^{\infty} \frac{(|z|\, q^n)^{k+1}}{(1-q)^{k+1}} \frac{(1-q)^{k+1} q^{k(k+1)/2}}{(q;q)_{k+1}} \\ &\le \sum_{k=0}^{\infty} \frac{(|z|\, q^n)^{k+1}}{(1-q)^{k+1}} \frac{1}{(k+1)!} \le \frac{|z|\, q^n \sqrt{e}}{1-q} < \frac{2\,|z|\, q^n}{1-q}. \end{aligned} \tag{15}$$

□

*Remark.* In this work, we either have a fixed $q$ with $0 < q < 1$, $q = \exp(-d\pi n^{-a})$ or $q = \exp\left(-\frac{d}{\log n}\right)$, with $0 < a < 1$ and $d > 0$. In these cases, the condition (3) is clearly satisfied for $n$ large.

The Dedekind $\eta(\tau)$ is defined as [23]

$$\eta(\tau) := e^{\pi i\tau/12} \prod_{k=1}^{\infty} (1 - e^{2\pi i k\tau}), \tag{16}$$

or

$$\eta(\tau) = q^{1/12}(q^2;q^2)_\infty, \tag{17}$$

for

$$q = e^{\pi i\tau}, \quad \Im(\tau) > 0. \tag{18}$$

It has the transformation formula

$$\eta\left(-\frac{1}{\tau}\right) = \sqrt{\frac{\tau}{i}}\eta(\tau). \tag{19}$$

The four Jacobi theta functions are defined as

$$\theta_1(v|\tau) := -i\sum_{k=-\infty}^{\infty}(-1)^k q^{(k+1/2)^2}e^{(2k+1)\pi iv}, \tag{20}$$

$$\theta_2(v|\tau) := \sum_{k=-\infty}^{\infty} q^{(k+1/2)^2}e^{(2k+1)\pi iv}, \tag{21}$$

$$\theta_3(v|\tau) := \sum_{k=-\infty}^{\infty} q^{k^2}e^{2k\pi iv}, \tag{22}$$

$$\theta_4(v|\tau) := \sum_{k=-\infty}^{\infty} (-1)^k q^{k^2}e^{2k\pi iv}, \tag{23}$$

where

$$q = e^{\pi i\tau}, \quad \Im(\tau) > 0. \tag{24}$$

For our convenience, we also use the following notations

$$\theta_\lambda(z;q) := \theta_\lambda(v|\tau), \quad z = e^{2\pi iv}, \quad q = e^{\pi i\tau} \tag{25}$$

for $\lambda = 1,2,3,4$. The Jacobi's triple product identities are

$$\theta_1(v|\tau) = 2q^{1/4}\sin\pi v(q^2;q^2)_\infty(q^2e^{2\pi iv};q^2)_\infty(q^2e^{-2\pi iv};q^2)_\infty, \tag{26}$$

$$\theta_2(v|\tau) = 2q^{1/4}\cos\pi v(q^2;q^2)_\infty(-q^2e^{2\pi iv};q^2)_\infty(-q^2e^{-2\pi iv};q^2)_\infty, \tag{27}$$

$$\theta_3(v|\tau) = (q^2;q^2)_\infty(-qe^{2\pi iv};q^2)_\infty(-qe^{-2\pi iv};q^2)_\infty, \tag{28}$$

$$\theta_4(v|\tau) = (q^2;q^2)_\infty(qe^{2\pi iv};q^2)_\infty(qe^{-2\pi iv};q^2)_\infty. \tag{29}$$

The Jacobi $\theta$ functions satisfy transformations

$$\theta_1\left(\frac{v}{\tau} \mid -\frac{1}{\tau}\right) = -i\sqrt{\frac{\tau}{i}}e^{\pi iv^2/\tau}\theta_1\left(v \mid \tau\right), \tag{30}$$

$$\theta_2\left(\frac{v}{\tau} \mid -\frac{1}{\tau}\right) = \sqrt{\frac{\tau}{i}}e^{\pi iv^2/\tau}\theta_4\left(v \mid \tau\right), \tag{31}$$

$$\theta_3\left(\frac{v}{\tau} \mid -\frac{1}{\tau}\right) = \sqrt{\frac{\tau}{i}}e^{\pi iv^2/\tau}\theta_3\left(v \mid \tau\right), \tag{32}$$

$$\theta_4\left(\frac{v}{\tau} \mid -\frac{1}{\tau}\right) = \sqrt{\frac{\tau}{i}}e^{\pi iv^2/\tau}\theta_2\left(v \mid \tau\right). \tag{33}$$

**Lemma 1.2.** *Let*

$$0 < a < 1, \quad n \in \mathbb{N}, \tag{34}$$

*and*

$$q = e^{-2\pi\gamma^{-1}n^{-a}}, \quad \gamma > 0, \tag{35}$$

*then,*

$$(q;q)_\infty = \sqrt{\gamma n^a}\exp\left(\pi n^{-a}/(12\gamma) - \pi\gamma n^a/12\right)\left\{1 + \mathcal{O}\left(e^{-2\pi\gamma n^a}\right)\right\}, \tag{36}$$

*and*

$$\frac{1}{(q;q)_\infty} = \frac{\exp\left(\pi\gamma n^a/12 - \pi n^{-a}/(12\gamma)\right)}{\sqrt{\gamma n^a}}\left\{1 + \mathcal{O}\left(e^{-2\pi\gamma n^a}\right)\right\} \tag{37}$$

*as $n \to \infty$.*

*Proof.* From formulas (15) and (18) we get

$$\begin{aligned}(q;q)_\infty &= \exp\left(\pi n^{-a}/(12\gamma)\right)\eta\left(n^{-a}\gamma^{-1}i\right)\\ &= \sqrt{\gamma n^a}\exp\left(\pi n^{-a}/(12\gamma)\right)\eta(\gamma n^a i)\\ &= \sqrt{\gamma n^a}\exp\left(\pi n^{-a}/(12\gamma)-\pi\gamma n^a/12\right)\prod_{k=1}^{\infty}(1-e^{-2\pi\gamma k n^a})\\ &= \sqrt{\gamma n^a}\exp\left(\pi n^{-a}/(12\gamma)-\pi\gamma n^a/12\right)\left\{1+\mathcal{O}\left(e^{-2\pi\gamma n^a}\right)\right\}\end{aligned} \tag{38}$$

and

$$\frac{1}{(q;q)_\infty} = \frac{\exp\left(\pi\gamma n^a/12-\pi n^{-a}/(12\gamma)\right)}{\sqrt{\gamma n^a}}\left\{1+\mathcal{O}\left(e^{-2\pi\gamma n^a}\right)\right\} \tag{39}$$

as $n\to\infty$. ☐

Similarly, we have:

**Lemma 1.3.** *Assume that*

$$q = \exp\left(-\frac{2\pi}{\gamma\log n}\right), \quad n\ge 2, \quad \gamma>0, \tag{40}$$

*then,*

$$(q;q)_\infty = n^{-\pi\gamma/12}\sqrt{\gamma\log n}\exp\left(\frac{\pi}{12\gamma\log n}\right)\left\{1+\mathcal{O}\left(n^{-2\pi\gamma}\right)\right\}, \tag{41}$$

*and*

$$\frac{1}{(q;q)_\infty} = \frac{n^{\pi\gamma/12}}{\sqrt{\gamma\log n}}\exp\left(-\frac{\pi}{12\gamma\log n}\right)\left\{1+\mathcal{O}\left(n^{-2\pi\gamma}\right)\right\} \tag{42}$$

*as* $n\to\infty$.

We will also make use of the trivial inequality

$$|e^z-1|\le |z|e^{|z|} \tag{43}$$

for any $z\in\mathbb{C}$.

### 1.2. Generalized Irrational Measure.

For an irrational number $\theta$, Chebyshev's theorem implies that for any real number $\beta$ , there exist infinitely many pairs of integers $n$ and $m$ with $n>0$ such that [11]

$$n\theta = m+\beta+\gamma_n \quad \text{with} \quad |\gamma_n|\le\frac{3}{n}. \tag{44}$$

Clearly, Chebyshev's theorem says that the arithmetic progression $\{n\theta\}_{n\in\mathbb{Z}}$ is ergodic in $\mathbb{R}$.

**Definition 1.4.** Given real numbers $\theta_j$ and $\beta_j$ for $j=1,...,N$, the generalized irrationality measure $\omega(\theta_1,\dots,\theta_N|\beta_1,\dots,\beta_N)$ of $\theta_1,\dots,\theta_N$ associated with $\beta_1,\dots,\beta_N$ is defined as the least upper bound of the set of real numbers $r$ such that there exist infinitely many positive integers $n$ and integers $m_1,\dots,m_N$ such that

$$|n\theta_j-\beta_j-m_j| < \frac{1}{n^{r-1}}. \tag{45}$$

for $j=1,\dots,N$.

**Proposition 1.5.** *Let $\theta$ be an irrational number, then for any real number $\beta$, its generalized irrational measure associated with $\beta$ is $\omega(\theta|\beta)\ge 2$ .*

*Proof.* The assertion is a direct consequence of the Chebyshev's theorem. □

Recall that the irrationality measure (or Liouville-Roth constant) $\mu(\theta)$ of a real number $\theta$ is defined as the least upper bound of the set of real numbers $r$ such that [28]

$$0 < |n\theta - m| < \frac{1}{n^{r-1}} \tag{46}$$

is satisfied by an infinite number of integer pairs $(n, m)$ with $n > 0$. It is clear that we have

$$\omega(\theta|0) = \mu(\theta). \tag{47}$$

A real algebraic number $\theta$ of degree $\ell$ if it is a root of an irreducible polynomial of degree $\ell$ with integer coefficients. Liouville's theorem in number theory says that for a real algebraic number $\theta$ of degree $\ell$, there exists a positive constant $K(\theta)$ such that for any integer $m$ and $n > 0$ we have

$$|n\theta - m| > \frac{K(\theta)}{n^{\ell-1}}. \tag{48}$$

A Liouville number is a real number $\theta$ such that for any positive integer $\ell$ there exist infinitely many integers $n$ and $m$ with $n > 1$ such that [28]

$$0 < |n\theta - m| < \frac{1}{n^{\ell-1}}, \tag{49}$$

It is well known that even though the set of all Liouville numbers is of Lebesgue measure zero, topologically, almost all real numbers are Liouville numbers.

**Proposition 1.6.** *The generalized irrational measure $\omega(\theta|\beta)$ has the following properties:*

1. *For any real algebraic number $\theta$ of degree $\ell$, one has $\omega(\theta|0) \le \ell$.*
2. *For a Liouville number $\theta$, one has $\omega(\theta|0) = \infty$.*

*Proof.* The first assertion follows from the definition 1.4 and (47) while the last assertion follows directly from the definitions of the generalized irrational measure and the Liouville numbers. □

It is clear that the quadratic irrationals $\theta$ such as $\sqrt{2}$ have the generalized irrational measure $\omega(\theta|0) = 2$ for any real number $\beta$.

For any real number $\theta$, we consider the set

$$\mathbb{S}(\theta) = \{\{n\theta\} : n \in \mathbb{N}\}. \tag{50}$$

Obviously, the set $\mathbb{S}(\theta)$ is a finite subset of $[0, 1)$ for $\theta \in \mathbb{Q}$. When $\theta \notin \mathbb{Q}$, the set $\mathbb{S}(\theta)$ is dense in $(0, 1)$, which is a consequence of Chebyshev's theorem.

**Lemma 1.7.** *We have the following:*

1. *Assume that $\theta > 0$ and $\theta \in \mathbb{Q}$. For any real number $\lambda$ with $\{\lambda\} \in \mathbb{S}(\theta)$, there exist infinitely many pair of integers $n$ and $m$ with $n > 0$ such that*

$$n\theta = m + \lambda, \quad m \in \mathbb{N}. \tag{51}$$

2. *Let $\theta$ be an irrational number, then for any fixed real number $\beta$ there is a positive number $\rho \ge 1$ such that*

$$n\theta = m + \beta + \gamma_n, \quad m \in \mathbb{Z}, \quad |\gamma_n| \le \frac{1}{n^\rho} \tag{52}$$

*holds for infinitely many positive integers $n$. Furthermore, $m \in \mathbb{N}$ if $\theta > 0$.*

(3) *Given two rational numbers $\tau, \theta$ with $\tau < 0$, there are $\lambda$, $\lambda_1$ with $\{\lambda\} \in \mathbb{S}(-\tau)$, $\{\lambda_1\} \in \mathbb{S}(\theta)$ and infinitely many integers $n$, $m$ with $n > 0$ such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}, \quad n\theta = m_1 + \lambda_1, \quad m_1 \in \mathbb{Z}. \tag{53}$$

(4) *Assume $\tau < 0$. If only one of $\tau$ and $\theta$ is rational, say, $\tau$ is rational and $\theta$ is irrational, then for any fixed real number $\beta$, there is a positive number $\rho \geq 1$ and some rational number $\lambda$ with $\{\lambda\} \in S(-\tau)$ such that*

$$n\theta = m_1 + \beta + b_n, \quad m_1 \in \mathbb{Z}, \quad |b_n| < \frac{1}{n^\rho} \tag{54}$$

*and*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N} \tag{55}$$

*hold for infinitely many positive integers $n$.*

(5) *If both $-\tau > 0$ and $\theta$ are irrational and there exist real numbers $\beta_1$, $\beta_2$ such that*

$$\omega(-\tau, \theta|\beta_1, \beta_2) > 1, \tag{56}$$

*then for any $\rho$ with*

$$0 < \rho < \omega(-\tau, \theta|\beta_1, \beta_2) - 1, \tag{57}$$

*there exist infinitely many integers $n$ and $m$ with $n > 0$ such that*

$$n\theta = m_1 + \beta_2 + b_n, \quad m_1 \in \mathbb{Z}, \quad |b_n| < \frac{1}{n^\rho}, \tag{58}$$

*and*

$$-n\tau = m + \beta_1 + a_n, \quad m \in \mathbb{N}, \quad |a_n| < \frac{1}{n^\rho}. \tag{59}$$

*Proof.* The first two cases are trivial, they are direct consequences of definition of $\omega(\theta|\beta)$ and the Chebyshev's theorem.

To prove the third assertion, we only need to consider the case that $\lambda, \lambda_1 \in (0, 1)$. Let $\tau, \theta \in \mathbb{Q}$, if $\theta$ or $\tau$ is an integer, then (53) reduces to (51). Assume that $\tau < 0$ and $\theta$ are not integers, let

$$-\tau = \frac{q_1}{p_1}, \quad p_1 \geq 1, \quad (p_1, q_1) = 1, \tag{60}$$

and

$$\theta = \frac{q_2}{p_2}, \quad p_2 \geq 1, \quad (p_2, q_2) = 1. \tag{61}$$

If

$$d = (p_1, p_2) = 1, \tag{62}$$

and for any

$$\lambda = \frac{v_1}{p_1} \in \mathbb{S}(-\tau), \quad \lambda_1 = \frac{v_2}{p_2} \in \mathbb{S}(\theta), \tag{63}$$

let $u_1$, $u_2$ be any two integers satisfying

$$v_1 \equiv q_1 u_1 (\mod p_1), \quad v_2 \equiv q_2 u_2 (\mod p_2), \tag{64}$$

then, by Chinese remainder theorem, there exist infinitely many positive integers $n, n_1, n_2$ such that

$$n = p_1 n_1 + u_1, \quad n = p_2 n_2 + u_2. \tag{65}$$

Hence,

$$-n\tau = n_1 q_1 + \frac{q_1 u_1}{p_1} = m + \lambda \tag{66}$$

and

$$n\theta = n_2 q_2 + \frac{q_2 u_2}{p_2} = m_1 + \lambda_1. \tag{67}$$

In the case that

$$d = (p_1, p_2) > 1, \tag{68}$$

we let

$$p_1' = \frac{p_1}{d}, \quad p_2' = \frac{p_2}{d}. \tag{69}$$

From the previous discussion we see that for

$$\lambda \in \left\{0, \frac{1}{p_1'}, \dots, \frac{p_1'-1}{p_1'}\right\} \subset \mathbb{S}(-\tau) \tag{70}$$

and

$$\lambda_1 \in \left\{0, \frac{1}{p_2'}, \dots, \frac{p_2'-1}{p_2'}\right\} \subset \mathbb{S}(\theta), \tag{71}$$

there exists infinitely many positive integers $n'$, $m'$ such that

$$-n'\tau = m + \lambda, \quad n'\theta = m_1 + \lambda_1, \quad m_1, m \in \mathbb{Z}, \tag{72}$$

where the new $n' = dn$ for each $n$ in (64). Given any two real numbers $\tau < 0$ and $\theta$, if just one of them is irrational, say, $\tau$ is rational and $\theta$ is irrational, then by (51), there are infinitely many positive integers $n$ satisfying equations of (53). Since $\mathbb{S}(-\tau)$ is finite set, then there exist some $\lambda \in \mathbb{S}(-\tau)$ and infinitely many positive integers $n$ satisfying all the equations of (53) and (54).

In the case when both $\tau < 0$ and $\theta$ are irrational, (57) and (58) follow from the definition of $\omega(-\tau, \theta|\beta_1, \beta_2)$. □

## 2. $q$-Exponential Function $E_q(z)$

The Euler's $q$-Exponential is defined by [6, 9, 13, 20]

$$E_q(z) := (-z; q)_\infty = \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{(q;q)_k} z^k, \quad z \in \mathbb{C}. \tag{73}$$

For any complex number $z$, we notice that

$$E_q((1-q)z) = \sum_{k=0}^{\infty} \frac{(1-q)^k}{(q;q)_k} q^{k(k-1)/2} z^k. \tag{74}$$

By applying (13) we get

$$\frac{(1-q)^k q^{k(k-1)/2}}{(q;q)_k} \le \frac{1}{k!} \tag{75}$$

for any nonnegative integer $k$. Hence by Lebesgue dominated theorem we have

$$\lim_{q\to 1} E_q((1-q)z) = \sum_{k=0}^{\infty} \frac{z^k}{k!} = e^z, \tag{76}$$

and this is the reason why $E_q(z)$ is called a $q$-exponential. Indeed, it is one of several $q$-analogues of $e^z$ in $q$-series. From (73) and (74), it is also clear that

$$|E_q((1-q)z)| \le e^{|z|}, \quad z \in \mathbb{C}. \tag{77}$$

From (72) we also have

$$|(E_q(z)| \le \sum_{k=0}^{\infty} \frac{q^{k/2}}{(q;q)_k} \left\{ q^{k^2/2} \left(\frac{|z|}{q}\right)^k \right\} \tag{78}$$

and for any nonzero complex number $z$, the terms

$$q^{k^2/2} \left(\frac{|z|}{q}\right)^k, \quad k = 0, 1, ... \tag{79}$$

are bounded by

$$\exp\left\{-\frac{\log^2 |z/q|}{2\log q}\right\}. \tag{80}$$

Then we have

$$|(E_q(z)| \le \exp\left\{-\frac{\log^2 |z/q|}{2\log q}\right\} \sum_{k=0}^{\infty} \frac{q^{k/2}}{(q;q)_k} \tag{81}$$

or

$$|(E_q(z)| \le \frac{\exp\left\{-\frac{\log^2 |z/q|}{2\log q}\right\}}{(\sqrt{q};q)_\infty}. \tag{82}$$

From Jacobi triple product formulas we obtain

$$\begin{aligned} |\theta_3(z;q)| &= \left|(q^2, -qz, -q/z; q^2)_\infty\right| \\ &\le \frac{(q^2;q^2)_\infty}{(q;q^2)_\infty^2} \exp\left\{-\frac{1}{4\log q}\left[\log^2(|z|/q) + \log^2(|z|q)\right]\right\}. \end{aligned} \tag{83}$$

Since

$$\log^2(|z|/q) + \log^2(|z|q) = 2\log^2|z| + 2\log^2 q, \tag{84}$$

then,

$$|\theta_3(z;q)| \le \frac{(q^2;q^2)_\infty}{(q;q^2)_\infty^2 \sqrt{q}} \exp\left\{-\frac{\log^2|z|}{2\log q}\right\}. \tag{85}$$

Similarly, we have

$$|\theta_4(z;q)| \quad \le \quad \frac{(q^2;q^2)_\infty}{(q;q^2)_\infty^2 \sqrt{q}} \exp\left\{-\frac{\log^2|z|}{2\log q}\right\}, \tag{86}$$

$$|\theta_1(z;q)| \quad \le \quad \frac{2\sqrt[4]{q}(q^2;q^2)_\infty \cosh(\frac{1}{2}\log|z|)}{(q;q^2)_\infty^2} \exp\left\{-\frac{\log^2|z|}{2\log q}\right\}, \tag{87}$$

$$|\theta_2(z;q)| \quad \le \quad \frac{2\sqrt[4]{q}(q^2;q^2)_\infty \cosh(\frac{1}{2}\log|z|)}{(q;q^2)_\infty^2} \exp\left\{-\frac{\log^2|z|}{2\log q}\right\}. \tag{88}$$

### 2.1. **Asymptotics for $E_q(z)$ .** Assume that

$$s=\tau+\frac{2\pi i\theta}{\log q},\quad \tau,\theta\in\mathbb{R}. \tag{89}$$

Then for any nonzero complex number $z$, we have

$$E_q(-q^{ns+1/2}z) \quad = \quad \sum_{k=0}^{\infty}\frac{q^{k^2/2}}{(q;q)_k}\left(-ze^{2\pi in\theta}q^{n\tau}\right)^k. \tag{90}$$

For the Euler $q$-Exponential $E_q(z)$ we have the following results:

**Theorem.** *For any nonzero complex number $z$, let*

$$j_n=\left\lfloor\frac{q^4\log n}{-\log q}\right\rfloor,\quad k_n=\left\lfloor\frac{-n\tau}{2}\right\rfloor, \tag{91}$$

*we have the following results for $E_q(z)$:*

(1) *If $\tau>0$, then we have*

$$E_q(-q^{ns+1/2}z)=1+r_{qe}(n|1), \tag{92}$$

*and*

$$|r_{qe}(n|1)|\le\frac{|z|q^{n\tau+1/2}}{1-q}\exp\left\{\frac{|z|q^{n\tau+1/2}}{1-q}\right\}. \tag{93}$$

(2) *Assume that $\tau=0$. If for some fixed real numbers $\beta$ and $\rho\ge 1$ there are infinitely many positive integers $n$ such that*

$$n\theta=m+\beta+b_n,\quad |b_n|<\frac{1}{n^\rho},\quad m\in\mathbb{Z}, \tag{94}$$

*then,*

$$E_q(-q^{ns+1/2}z)=E_q(-zq^{1/2}e^{2\pi i\beta};q)_\infty+r_{qe}(n|3), \tag{95}$$

*and*

$$|r_{qe}(n|3)|\le 24\exp\left(\frac{|z|\sqrt{q}}{1-q}\right)\left\{\frac{\log n}{n^\rho}+\frac{1}{j_n!}\left(\frac{|z|\sqrt{q}}{1-q}\right)^{j_n}\right\} \tag{96}$$

*for $n$ sufficiently large.*

(3) *Assume that $\tau<0$. If for some fixed real numbers $\lambda$ and $\lambda_1$ there are infinitely many positive integers $n$ such that*

$$-n\tau=m+\lambda,\quad m\in\mathbb{N},\quad n\theta=m_1+\lambda_1,\quad m_1\in\mathbb{Z}. \tag{97}$$

*Then*

$$\frac{E_q(-q^{ns+1/2}z)(q;q)_\infty}{(-ze^{2\pi in\theta})^m q^{m(n\tau+m/2)}}=\theta_4\left(z^{-1}q^\lambda e^{-2\pi i\lambda_1};\sqrt{q}\right)+r_{qe}(n|4), \tag{98}$$

*and*

$$|r_{qe}(n|4)|\le 4\theta_3\left(|z|^{-1}q^\lambda;\sqrt{q}\right)\left\{\frac{q^{k_n}}{1-q}+\frac{q^{k_n^2/2+\lambda k_n}}{|z|^{k_n}}\right\} \tag{99}$$

*for $n$ sufficiently large.*

(4) *Assume that $\tau < 0$. If for some fixed real numbers $\beta$, $\lambda$, $\rho \geq 1$ there exist infinitely many positive integers $n$ such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}, \tag{100}$$

*and*

$$n\theta = m_1 + \beta + b_n, \quad |b_n| < \frac{1}{n^\rho}, \quad m_1 \in \mathbb{Z}, \tag{101}$$

*then,*

$$\frac{E_q(-q^{ns+1/2}z)(q;q)_\infty}{(-ze^{2\pi i n\theta})^m q^{m(n\tau+m/2)}} = \theta_4\left(z^{-1}q^{\lambda}e^{-2\pi i\beta};\sqrt{q}\right) + r_{qe}(n|5), \tag{102}$$

*and*

$$|r_{qe}(n|5)| \leq 48\theta_3\left(|z|^{-1}q^{\lambda};\sqrt{q}\right)\left\{\frac{q^{j_n^2/2+\lambda j_n}}{|z|^{j_n}} + |z|^{j_n}\, q^{j_n^2/2-\lambda j_n} + \frac{q^{k_n}}{1-q} + \frac{\log n}{n^\rho}\right\} \tag{103}$$

*for $n$ sufficiently large.*

(5) *Assume that $\tau < 0$. If for some fixed real numbers $\beta$, $\lambda$, $\rho \geq 1$ there are infinitely many positive integers $n$ such that*

$$-n\tau = m + \beta + a_n, \quad |a_n| < \frac{1}{n^\rho}, \quad m \in \mathbb{N}, \quad n\theta = m_1 + \lambda, \quad m_1 \in \mathbb{Z}, \tag{104}$$

*then,*

$$\frac{E_q(-q^{ns+1/2}z)(q;q)_\infty}{(-ze^{2\pi i n\theta})^m q^{m(n\tau+m/2)}} = \theta_4\left(z^{-1}q^{\beta}e^{-2\pi i\lambda};\sqrt{q}\right) + r_{qe}(n|6), \tag{105}$$

*and*

$$|r_{qe}(n|6)| \leq 6\theta_3(|z|^{-1}q^{\beta};\sqrt{q})\left\{\frac{\log n}{n^\rho} + \frac{q^{k_n}}{1-q} + |z|^{j_n}\, q^{j_n^2/2-(\beta+1)j_n} + \frac{q^{j_n^2/2+(\beta-1)j_n}}{|z|^{j_n}}\right\} \tag{106}$$

*for $n$ sufficiently large.*

(6) *Assume that $\tau < 0$. If for some fixed real numbers $\beta_1$, $\beta_2$ and $\rho > 0$ there exist infinitely many positive integers $n$ such that*

$$-n\tau = m + \beta_1 + a_n, \quad |a_n| < \frac{1}{n^\rho}, \quad m \in \mathbb{N}, \tag{107}$$

*and*

$$n\theta = m_1 + \beta_2 + b_n, \quad |b_n| < \frac{1}{n^\rho}, \quad m_1 \in \mathbb{Z}. \tag{108}$$

*Then,*

$$\frac{E_q(-q^{ns+1/2}z)(q;q)_\infty}{(-ze^{2\pi i n\theta})^m q^{m(n\tau+m/2)}} = \theta_4\left(z^{-1}q^{\beta_1}e^{-2\pi i\beta_2};\sqrt{q}\right) + r_{qe}(n|7), \tag{109}$$

*and*

$$|r_{qe}(n|7)| \leq 54\theta_3(|z|^{-1}q^{\beta_1};\sqrt{q})\left\{\frac{\log n}{n^\rho} + \frac{q^{k_n}}{1-q} + q^{j_n^2/2-(\beta_1+1)j_n}|z|^{j_n} + \frac{q^{j_n^2/2+(\beta_1-1)j_n}}{|z|^{j_n}}\right\} \tag{110}$$

*for $n$ sufficiently large.*

*Remark* 2.1. If only one of $\tau$ and $\theta$ is an irrational number, say $\tau$, then the real number $\beta$ could be fixed as 0. If $\tau$ is an algebraic of degree $\ell$, then the order of error term can be no better than $\mathcal{O}(n^{1-\ell})$, while if the number is a Liouville number, then, the order is better than $\mathcal{O}(n^{-r})$ for any real number $r$. In case of an arbitrary irrational number $\tau$ and an arbitrary real number $\beta$, the error term is $\mathcal{O}(n^{-1+\epsilon})$ for any small $\epsilon > 0$ by the Chebyshev's theorem. This remark applies to all the asymptotics in this work.

In the following corollaries we assume that

$$z := e^{2\pi u}, \quad u \in \mathbb{R}. \tag{111}$$

**Corollary 2.2.** *Assume that*

$$q = \exp(-2n^{-a}\pi), \quad 0 < a < \frac{1}{2}, \quad n \in \mathbb{N}, \tag{112}$$

*we have the following results for* $E_q(z)$*:*

1. *Assume that* $\tau > 0$*, we have*

$$E_q(-\exp 2\pi(u + n^{1-a}\tau + n^{-a}/2)) = 1 + \mathcal{O}\left(n^a e^{-2\pi\tau n^{1-a}}\right) \tag{113}$$

   *for* $n$ *sufficiently large.*

2. *Assume that* $\tau < 0$*, if for some fixed real number* $\lambda$ *there are infinitely many positive integer* $n$ *such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}, \tag{114}$$

   *then,*

$$\begin{aligned} &E_q(-\exp 2\pi(u + \tau n^{1-a} + n^{-a}/2)) \\ &= \frac{2\exp\left(\pi n^{-a}(n^a u - n\tau)^2\right)\left\{\cos\pi(n^a u + \lambda) + \mathcal{O}\left(e^{-2\pi n^a}\right)\right\}}{(-1)^{\lambda + n\tau}\exp\left(\pi n^a/6 + \pi n^{-a}/12\right)} \end{aligned} \tag{115}$$

   *for* $n$ *sufficiently large.*

Let

$$q = \exp(-\frac{2\pi}{\gamma\log n}), \quad n \geq 2, \quad \gamma > 0, \tag{116}$$

then we have the following:

**Corollary 2.3.** *we have the following results for* $E_q(z)$*:*

1. *Assume that* $\tau > 0$*, we have*

$$E_q\left(-\exp 2\pi\left(u - \frac{n\tau + 1/2}{\gamma\log n}\right)\right) = 1 + \mathcal{O}\left(\log n \exp\left(-\frac{2n\pi\tau}{\gamma\log n}\right)\right) \tag{117}$$

   *for* $n$ *sufficiently large.*

2. *Assume that* $\tau < 0$ *and for some fixed real number* $\lambda$ *there are infinitely many positive integer* $n$ *such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}, \tag{118}$$

*then,*

$$\begin{aligned} E_q\left(-\exp 2\pi\left(u-\frac{n\tau+1/2}{\gamma\log n}\right)\right) &\\ = (-1)^{\tau n+\lambda}\exp\left(\frac{\pi(\gamma u\log n-n\tau)^2}{\gamma\log n}-\frac{\pi}{12\gamma\log n}\right) &\\ \times 2n^{-\gamma\pi/6}\left\{\cos\pi(\gamma u\log n+\lambda)+\mathcal{O}(n^{-2\pi\gamma})\right\} & \end{aligned} \tag{119}$$

*for $n$ sufficiently large.*

(3) *Assume that $\tau<0$, if for some fixed real numbers $\beta$, $\rho\geq 1$ and $\lambda$ there exist infinitely many positive integers $n$ such that*

$$-n\tau=m+\beta+a_n,\quad |a_n|<\frac{1}{n^{\rho}},\quad m\in\mathbb{N}, \tag{120}$$

*then,*

$$\begin{aligned} E_q\left(-\exp 2\pi\left(u-\frac{n\tau+1/2}{\gamma\log n}\right)\right) &\\ =\exp\left(\frac{\pi(\gamma u\log n-n\tau)^2}{\gamma\log n}-\frac{\pi}{12\gamma\log n}\right) &\\ \times\frac{2\left\{\cos\pi(\gamma u\log n+\beta)+\mathcal{O}(n^{-8\rho/9}\log n)\right\}}{(-1)^m n^{2\rho/27}} & \end{aligned} \tag{121}$$

*for $n$ sufficiently large, where*

$$\gamma=\frac{4\rho}{9\pi}. \tag{122}$$

### 2.2. **Proofs for Theorem 2.1.**

Assume that $\tau<0$ and

$$-n\tau=m+c_n,\quad m\in\mathbb{N},\quad n\theta=m_1+d_n,\quad m_1\in\mathbb{Z}, \tag{123}$$

then,

$$\begin{aligned} E(-q^{ns+1/2}z) &=\sum_{k=0}^{\infty}\frac{q^{k^2/2}}{(q;q)_k}\left(-ze^{2\pi in\theta}q^{n\tau}\right)^k\\ &=\sum_{k=0}^{m}\frac{q^{k^2/2}}{(q;q)_k}\left(-ze^{2\pi in\theta}q^{n\tau}\right)^k+\sum_{k=m+1}^{\infty}\frac{q^{k^2/2}}{(q;q)_k}\left(-ze^{2\pi in\theta}q^{n\tau}\right)^k\\ &=s_1+s_2. \end{aligned} \tag{124}$$

Reverse summation order in $s_1$,

$$\frac{(q;q)_\infty s_1}{(-ze^{2\pi in\theta})^m q^{m(n\tau+m/2)}}=\sum_{k=0}^{m}q^{k^2/2}(-q^{c_n}z^{-1}e^{-2\pi id_n})^k e(k,n) \tag{125}$$

with

$$e(k,n)=\frac{(q;q)_\infty}{(q;q)_{m-k}}, \tag{126}$$

then,

$$|e(k,n)|\leq 1 \tag{127}$$

for $0 \le k \le m$. An application of Lemma 1.1 gives

$$|e(k,n)-1| = |r_1(q;m-k)| \le \frac{2q^{2+k_n}}{1-q}. \tag{128}$$

for $0 \le k \le k_n - 1$ and $n$ sufficiently large.

We shift the summation index from $k$ to $k+m$ in $s_2$,

$$\frac{(q;q)_\infty s_2}{(-ze^{2\pi in\theta})^m q^{m(n\tau+m/2)}} = \sum_{k=1}^{\infty} q^{k^2/2}(-zq^{-c_n}e^{2\pi id_n})^k f(k,n), \tag{129}$$

with

$$f(k,n) = \frac{(q;q)_\infty}{(q;q)_{m+k}}, \tag{130}$$

then,

$$|f(k,n)| \le 1, \tag{131}$$

and

$$|f(k,n)-1| = |r_1(q;m+k)| \le \frac{2q^{k_n+2}}{1-q} \tag{132}$$

for $k \in \mathbb{N}$ and $n$ sufficiently large.

2.2.1. *Proof for case 1.* If we write

$$E_q(-q^{ns+1/2}z) = 1 + r_{qe}(n|1), \tag{133}$$

then,

$$r_{qe}(n|1) = \sum_{k=1}^{\infty} \frac{q^{k^2/2}}{(q;q)_k}\left(-ze^{2\pi in\theta}q^{n\tau}\right)^k. \tag{134}$$

Hence,

$$\begin{aligned} |r_{qe}(n|1)| &\le \sum_{k=1}^{\infty} \frac{q^{k^2/2}}{(q;q)_k}(|z|q^{n\tau})^k \le \sum_{k=1}^{\infty} \frac{(|z|q^{n\tau+1/2})^k}{k!(1-q)^k}\frac{(1-q)^k k! q^{k(k-1)/2}}{(q;q)_k} \\ &\le \frac{|z|q^{n\tau+1/2}}{1-q}\sum_{k=0}^{\infty}\frac{(|z|q^{n\tau+1/2})^k}{k!(1-q)^k} \le \frac{|z|q^{n\tau+1/2}}{1-q}\exp\left\{\frac{|z|q^{n\tau+1/2}}{1-q}\right\} \end{aligned} \tag{135}$$

by an application of (13).

2.2.2. *Proof for case 2.* It is clear that

$$1 < j_n < \frac{n^\rho}{2\pi} \tag{136}$$

for $n$ sufficiently large, then,

$$
\begin{aligned}
E(-q^{ns+1/2}z) &= \sum_{k=0}^{\infty} \frac{q^{k^2/2}}{(q;q)_k} \left(-ze^{2\pi i\beta}\right)^k e^{2\pi ikb_n} \\
&= \sum_{k=0}^{\infty} \frac{q^{k^2/2}}{(q;q)_k} \left(-ze^{2\pi i\beta}\right)^k - \sum_{k=j_n}^{\infty} \frac{q^{k^2/2}}{(q;q)_k} \left(-ze^{2\pi i\beta}\right)^k \\
&+ \sum_{k=0}^{j_n-1} \frac{q^{k^2/2}}{(q;q)_k} \left(-ze^{2\pi i\beta}\right)^k \left\{e^{2\pi ikb_n}-1\right\} + \sum_{k=j_n}^{\infty} \frac{q^{k^2/2}}{(q;q)_k} \left(-ze^{2\pi i\beta}\right)^k e^{2\pi ikb_n} \\
&= E_q(-zq^{1/2}e^{2\pi i\beta}) + s_1 + s_2 + s_3.
\end{aligned} \tag{137}
$$

Then,

$$
\begin{aligned}
|s_1+s_3| &\le 2\sum_{k=j_n}^{\infty} \frac{q^{k^2/2}}{(q;q)_k}|z|^k \le 2\sum_{k=j_n}^{\infty} \frac{1}{k!}\left(\frac{|z|\sqrt{q}}{1-q}\right)^k \\
&\le \frac{\left(\frac{|z|\sqrt{q}}{1-q}\right)^{j_n}}{j_n!} \sum_{k=0}^{\infty} \frac{\left(\frac{|z|\sqrt{q}}{1-q}\right)^k}{k!} \le \frac{2\exp\left(\frac{|z|\sqrt{q}}{1-q}\right)}{j_n!}\left(\frac{|z|\sqrt{q}}{1-q}\right)^{j_n}.
\end{aligned} \tag{138}
$$

For $n$ sufficiently large,

$$
|s_2| \le 2\pi j_n|b_n|e^{2\pi j_n|b_n|}\sum_{k=0}^{\infty}\frac{q^{k^2/2}|z|^k}{(q;q)_k} \le 24\exp\left(\frac{|z|\sqrt{q}}{1-q}\right)\frac{\log n}{n^\rho}. \tag{139}
$$

Hence we have proved

$$
E_q(-q^{ns+1/2}z) = E_q(-zq^{1/2}e^{2\pi i\beta}) + r_{qe}(n|3), \tag{140}
$$

where

$$
r_{qe}(n|3) = s_1 + s_2 + s_3, \tag{141}
$$

and

$$
|r_{qe}(n|3)| \le 24\exp\left(\frac{|z|\sqrt{q}}{1-q}\right)\left\{\frac{\log n}{n^\rho} + \frac{1}{j_n!}\left(\frac{|z|\sqrt{q}}{1-q}\right)^{j_n}\right\} \tag{142}
$$

for $n$ sufficiently large.

2.2.3. *Proof for case 4.* Notice that

$$
\begin{aligned}
\frac{(q;q)_\infty s_1}{(-ze^{2\pi in\theta})^m q^{m(n\tau+m/2)}} &= \sum_{k=0}^{m} q^{k^2/2}(-q^\lambda z^{-1}e^{-2\pi i\lambda_1})^k e(k,n) \\
&= \sum_{k=0}^{\infty} q^{k^2/2}(-q^\lambda z^{-1}e^{-2\pi i\lambda_1})^k - \sum_{k=k_n}^{\infty} q^{k^2/2}(-q^\lambda z^{-1}e^{-2\pi i\lambda_1})^k \\
&+ \sum_{k=0}^{k_n-1} q^{k^2/2}(-q^\lambda z^{-1}e^{-2\pi i\lambda_1})^k\left\{e(k,n)-1\right\} + \sum_{k=k_n}^{m} q^{k^2/2}(-q^\lambda z^{-1}e^{-2\pi i\lambda_1})^k e(k,n) \\
&= \sum_{k=0}^{\infty} q^{k^2/2}(-q^\lambda z^{-1}e^{-2\pi i\lambda_1})^k + s_{11} + s_{12} + s_{13}.
\end{aligned} \tag{143}
$$

Thus,

$$|s_{11}+s_{13}| \le 2\sum_{k=k_n}^{\infty} q^{k^2/2}\left(\frac{q^{\lambda}}{|z|}\right)^{k} \le 2\frac{q^{k_n^2/2+\lambda k_n}}{|z|^{k_n}}\sum_{k=0}^{\infty} q^{k^2/2}\left(\frac{q^{\lambda+k_n}}{|z|}\right)^{k} \le 2\theta_3\left(|z|^{-1}q^{\lambda};\sqrt{q}\right)\frac{q^{k_n^2/2+\lambda k_n}}{|z|^{k_n}}, \tag{144}$$

and for $n$ sufficiently large,

$$|s_{12}| \le \frac{2q^{k_n}}{1-q}\sum_{k=0}^{\infty} q^{k^2/2}\left(\frac{q^{\lambda}}{|z|}\right)^{k} \le \frac{2\theta_3\left(|z|^{-1}q^{\lambda};\sqrt{q}\right)}{1-q}q^{k_n}. \tag{145}$$

Let

$$r_1(n) = s_{11}+s_{12}+s_{13}, \tag{146}$$

then,

$$\frac{(q;q)_\infty s_1}{(-ze^{2\pi in\theta})^m q^{m(n\tau+m/2)}} = \sum_{k=0}^{\infty} q^{k^2/2}(-q^{\lambda}z^{-1}e^{-2\pi i\lambda_1})^k + r_1(n), \tag{147}$$

and

$$|r_1(n)| \le 2\theta_3\left(|z|^{-1}q^{\lambda};\sqrt{q}\right)\left\{\frac{q^{k_n}}{1-q}+\frac{q^{k_n^2/2+\lambda k_n}}{|z|^{k_n}}\right\} \tag{148}$$

for $n$ sufficiently large.

In the second sum,

$$\begin{aligned}\frac{(q;q)_\infty s_2}{(-ze^{2\pi in\theta})^m q^{m(n\tau+m/2)}} &= \sum_{k=1}^{\infty} q^{k^2/2}(-zq^{-\lambda}e^{2\pi i\lambda_1})^k f(k,n) \\ &= \sum_{k=1}^{\infty} q^{k^2/2}(-zq^{-\lambda}e^{2\pi i\lambda_1})^k + \sum_{k=1}^{\infty} q^{k^2/2}(-zq^{-\lambda}e^{2\pi i\lambda_1})^k\left\{f(k,n)-1\right\} \\ &= \sum_{k=-\infty}^{-1} q^{k^2/2}(-q^{\lambda}z^{-1}e^{-2\pi i\lambda_1})^k + r_2(n),\end{aligned} \tag{149}$$

and

$$|r_2(n)| \le \frac{2q^{k_n}}{1-q}\sum_{k=1}^{\infty} q^{k^2/2}(q^{-\lambda}|z|)^k \le \frac{2\theta_3\left(|z|^{-1}q^{\lambda};\sqrt{q}\right)}{1-q}q^{k_n} \tag{150}$$

for $n$ sufficiently large. Hence,

$$\frac{E_q(-q^{ns+1/2}z)(q;q)_\infty}{(-ze^{2\pi in\theta})^m q^{m(n\tau+m/2)}} = \theta_4\left(z^{-1}q^{\lambda}e^{-2\pi i\lambda_1};\sqrt{q}\right) + r_{qe}(n|4), \tag{151}$$

with

$$|r_{qe}(n|4)| \le 4\theta_3\left(|z|^{-1}q^{\lambda};\sqrt{q}\right)\left\{\frac{q^{k_n}}{1-q}+\frac{q^{k_n^2/2+\lambda k_n}}{|z|^{k_n}}\right\} \tag{152}$$

for $n$ sufficiently large.

2.2.4. *Proof for case 5.* In this case we have

$$
\begin{aligned}
&\frac{(q;q)_\infty s_1}{(-ze^{2\pi in\theta})^m q^{m(n\tau+m/2)}} = \sum_{k=0}^{m} q^{k^2/2}(-q^\lambda z^{-1}e^{-2\pi i\beta})^k e^{-2\pi ikb_n} e(k,n) \\
&= \sum_{k=0}^{\infty} q^{k^2/2}(-q^\lambda z^{-1}e^{-2\pi i\beta})^k - \sum_{k=j_n}^{\infty} q^{k^2/2}(-q^\lambda z^{-1}e^{-2\pi i\beta})^k \\
&+ \sum_{k=0}^{j_n-1} q^{k^2/2}(-q^\lambda z^{-1}e^{-2\pi i\beta})^k \left\{e(k,n)-1\right\} + \sum_{k=0}^{j_n-1} q^{k^2/2}(-q^\lambda z^{-1}e^{-2\pi i\beta})^k e(k,n) \left\{e^{-2\pi ikb_n}-1\right\} \\
&+ \sum_{k=j_n}^{m} q^{k^2/2}(-q^\lambda z^{-1}e^{-2\pi i\beta})^k e^{-2\pi ikb_n} e(k,n) \\
&= \sum_{k=0}^{\infty} q^{k^2/2}(-q^\lambda z^{-1}e^{-2\pi i\beta})^k + s_{11}+s_{12}+s_{13}+s_{14}.
\end{aligned} \tag{153}
$$

Then,

$$
\begin{aligned}
|s_{11}+s_{14}| &\le 2\sum_{k=j_n}^{\infty} q^{k^2/2}\left(\frac{q^\lambda}{|z|}\right)^k \le 2\frac{q^{j_n^2/2+\lambda j_n}}{|z|^{j_n}}\sum_{k=0}^{\infty} q^{k^2/2}\left(\frac{q^{\lambda+j_n}}{|z|}\right)^k \\
&\le 2\theta_3\left(|z|^{-1}q^\lambda;\sqrt{q}\right)\frac{q^{j_n^2/2+\lambda j_n}}{|z|^{j_n}}.
\end{aligned} \tag{154}
$$

We also have

$$
|s_{13}| \le 2\pi j_n|b_n|e^{2\pi j_n|b_n|}\sum_{k=0}^{\infty} q^{k^2/2}\left(\frac{q^\lambda}{|z|}\right)^k \le 24\theta_3\left(|z|^{-1}q^\lambda;\sqrt{q}\right)\frac{\log n}{n^\rho}, \tag{155}
$$

and

$$
|s_{12}| \le \frac{2q^{k_n}}{1-q}\theta_3\left(|z|^{-1}q^\lambda;\sqrt{q}\right) \tag{156}
$$

for $n$ sufficiently large.

Let

$$
r_1(n) = s_{11}+s_{12}+s_{13}+s_{14}, \tag{157}
$$

then

$$
\frac{(q;q)_\infty s_1}{(-ze^{2\pi in\theta})^m q^{m(n\tau+m/2)}} = \sum_{k=0}^{\infty} q^{k^2/2}(-z^{-1}q^\lambda e^{-2\pi i\beta})^k + r_1(n), \tag{158}
$$

and

$$
|r_1(n)| \le 24\theta_3\left(|z|^{-1}q^\lambda;\sqrt{q}\right)\left\{\frac{q^{j_n^2/2+\lambda j_n}}{|z|^{j_n}} + \frac{q^{k_n}}{1-q} + \frac{\log n}{n^\rho}\right\} \tag{159}
$$

for $n$ sufficiently large.

Similarly,

$$\begin{aligned}\frac{(q;q)_\infty s_2}{(-ze^{2\pi in\theta})^m q^{m(n\tau+m/2)}} &= \sum_{k=1}^{\infty} q^{k^2/2}(-zq^{-\lambda}e^{2\pi i\beta})^k f(k,n)e^{2\pi ikb_n} \\ &= \sum_{k=1}^{\infty} q^{k^2/2}(-zq^{-\lambda}e^{2\pi i\beta})^k - \sum_{k=j_n}^{\infty} q^{k^2/2}(-zq^{-\lambda}e^{2\pi i\beta})^k \\ &+ \sum_{k=1}^{j_n-1} q^{k^2/2}(-zq^{-\lambda}e^{2\pi i\beta})^k \{f(k,n)-1\} + \sum_{k=1}^{j_n-1} q^{k^2/2}(-zq^{-\lambda}e^{2\pi i\beta})^k f(k,n)\left\{e^{2\pi ikb_n}-1\right\} \\ &+ \sum_{k=j_n}^{\infty} q^{k^2/2}(-zq^{-\lambda}e^{2\pi i\beta})^k f(k,n)e^{2\pi ikb_n} \\ &= \sum_{k=-\infty}^{-1} q^{k^2/2}(-z^{-1}q^{\lambda}e^{-2\pi i\beta})^k + s_{21}+s_{22}+s_{23}+s_{24}.\end{aligned} \tag{160}$$

Then,

$$\begin{aligned}|s_{21}+s_{24}| &\le 2\sum_{k=j_n}^{\infty} q^{k^2/2}(|z|q^{-\lambda})^k \le 2\,|z|^{j_n}\,q^{j_n^2/2-\lambda j_n}\sum_{k=0}^{\infty} q^{k^2/2}(|z|q^{-\lambda+j_n})^k \\ &\le 2\theta_3\left(|z|^{-1}q^{\lambda};\sqrt{q}\right)|z|^{j_n}\,q^{j_n^2/2-\lambda j_n}.\end{aligned} \tag{161}$$

For sufficiently large $n$ we have

$$|s_{22}| \le \frac{2q^{k_n}}{1-q}\sum_{k=1}^{\infty} q^{k^2/2}(|z|q^{-\lambda})^k \le \frac{2q^{k_n}}{1-q}\theta_3\left(|z|^{-1}q^{\lambda};\sqrt{q}\right), \tag{162}$$

and

$$|s_{23}| \le 2\pi j_n|b_n|e^{2\pi j_n|b_n|}\sum_{k=1}^{\infty} q^{k^2/2}(|z|q^{-\lambda})^k \le 24\theta_3\left(|z|^{-1}q^{\lambda};\sqrt{q}\right)\frac{\log n}{n^{\rho}} \tag{163}$$

for $n$ sufficiently large.

Let

$$r_2(n) = s_{21}+s_{22}+s_{23}+s_{24}, \tag{164}$$

then,

$$\frac{(q;q)_\infty s_2}{(-ze^{2\pi in\theta})^m q^{m(n\tau+m/2)}} = \sum_{k=-\infty}^{-1} q^{k^2/2}(-z^{-1}q^{\lambda}e^{-2\pi i\beta})^k + r_2(n), \tag{165}$$

with

$$|r_2(n)| \le 24\theta_3\left(|z|^{-1}q^{\lambda};\sqrt{q}\right)\left\{\frac{\log n}{n^{\rho}} + \frac{q^{k_n}}{1-q} + |z|^{j_n}\,q^{j_n^2/2-\lambda j_n}\right\} \tag{166}$$

for sufficiently large $n$.

Therefore,

$$\frac{E_q(-q^{ns+1/2}z)(q;q)_\infty}{(-ze^{2\pi in\theta})^m q^{m(n\tau+m/2)}} = \theta_4\left(z^{-1}q^{\lambda}e^{-2\pi i\beta};\sqrt{q}\right) + r_{qe}(n|5) \tag{167}$$

with

$$|r_{qe}(n|5)| \le 48\theta_3\left(|z|^{-1}q^{\lambda};\sqrt{q}\right)\left\{\frac{q^{j_n^2/2+\lambda j_n}}{|z|^{j_n}} + |z|^{j_n}\,q^{j_n^2/2-\lambda j_n} + \frac{q^{k_n}}{1-q} + \frac{\log n}{n^{\rho}}\right\} \tag{168}$$

for $n$ sufficiently large.

2.2.5. *Proof for case 6.* Notice that

$$\begin{aligned} &\frac{(q;q)_\infty s_1}{(-ze^{2\pi i n\theta})^m q^{m(n\tau+m/2)}} = \sum_{k=0}^{m} q^{k^2/2}(-q^{\beta}z^{-1}e^{-2\pi i\lambda})^k q^{ka_n} e(k,n) \\ &= \sum_{k=0}^{\infty} q^{k^2/2}(-q^{\beta}z^{-1}e^{-2\pi i\lambda})^k - \sum_{k=j_n}^{\infty} q^{k^2/2}(-q^{\beta}z^{-1}e^{-2\pi i\lambda})^k \\ &+ \sum_{k=0}^{j_n-1} q^{k^2/2}(-q^{\beta}z^{-1}e^{-2\pi i\lambda})^k\left\{e(k,n)-1\right\} + \sum_{k=0}^{j_n-1} q^{k^2/2}(-q^{\beta}z^{-1}e^{-2\pi i\lambda})^k e(k,n)\left\{q^{ka_n}-1\right\} \\ &+ \sum_{k=j_n}^{m} q^{k^2/2}(-q^{\beta}z^{-1}e^{-2\pi i\lambda})^k q^{ka_n} e(k,n) = \sum_{k=0}^{\infty} q^{k^2/2}(-q^{\beta}z^{-1}e^{-2\pi i\lambda})^k + s_{11}+s_{12}+s_{13}+s_{14}. \end{aligned} \tag{169}$$

Since

$$|q^{ka_n}-1| \le j_n|a_n|\log q^{-1}e^{j_n|a_n|\log q^{-1}} \le \frac{3\log n}{n^{\rho}}, \tag{170}$$

and

$$|q^{ka_n}| \le e^{j_n|a_n|\log q^{-1}} < 3 \tag{171}$$

for $0 \le k \le j_n-1$ and

$$|q^{ka_n}| \le q^{-2k} \tag{172}$$

for any $k \ge 0$, then,

$$\begin{aligned} |s_{11}+s_{14}| &\le 2\sum_{k=j_n}^{\infty} q^{k^2/2}(q^{\beta-1}|z|^{-1})^k \le \frac{2q^{j_n^2/2+(\beta-1)j_n}}{|z|^{j_n}}\sum_{k=0}^{\infty} q^{k^2/2}(q^{\beta-1+j_n}|z|^{-1})^k \\ &\le 2\theta_3\left(|z|^{-1}q^{\beta};\sqrt{q}\right)\frac{q^{j_n^2/2+(\beta-1)j_n}}{|z|^{j_n}}, \end{aligned} \tag{173}$$

and

$$|s_{12}| \le \frac{2q^{k_n}}{1-q}\sum_{k=0}^{\infty} q^{k^2/2}(q^{\beta}|z|^{-1})^k \le 2\theta_3(|z|^{-1}q^{\beta};\sqrt{q})\frac{q^{k_n}}{1-q}, \tag{174}$$

$$|s_{13}| \le \frac{3\log n}{n^{\rho}}\sum_{k=0}^{\infty} q^{k^2/2}(q^{\beta}|z|^{-1})^k \le 3\theta_3(|z|^{-1}q^{\beta};\sqrt{q})\frac{\log n}{n^{\rho}} \tag{175}$$

for $n$ sufficiently large. Hence,

$$\frac{(q;q)_\infty s_1}{(-ze^{2\pi i n\theta})^m q^{m(n\tau+m/2)}} = \sum_{k=0}^{\infty} q^{k^2/2}(-q^\beta z^{-1}e^{-2\pi i\lambda})^k + r_1(n), \tag{176}$$

where

$$r_1(n) = s_{11} + s_{12} + s_{13} + s_{14}, \tag{177}$$

and

$$|r_1(n)| \le 3\theta_3(|z|^{-1}q^\beta;\sqrt{q})\left\{\frac{q^{k_n}}{1-q} + \frac{\log n}{n^\rho} + \frac{q^{j_n^2/2+(\beta-1)j_n}}{|z|^{j_n}}\right\} \tag{178}$$

for $n$ sufficiently large. Similarly,

(179)

$$\frac{(q;q)_\infty s_2}{(-ze^{2\pi i n\theta})^m q^{m(n\tau+m/2)}} = \sum_{k=1}^{\infty} q^{k^2/2}(-zq^{-\beta}e^{2\pi i\lambda})^k f(k,n) q^{-ka_n}$$

$$= \sum_{k=-1}^{-\infty} q^{k^2/2}(-q^\beta z^{-1}e^{-2\pi i\lambda})^k - \sum_{k=j_n}^{\infty} q^{k^2/2}(-zq^{-\beta}e^{2\pi i\lambda})^k$$

$$+ \sum_{k=1}^{j_n-1} q^{k^2/2}(-zq^{-\beta}e^{2\pi i\lambda})^k\left\{f(k,n)-1\right\} + \sum_{k=1}^{j_n-1} q^{k^2/2}(-zq^{-\beta}e^{2\pi i\lambda})^k f(k,n)\left\{q^{-ka_n}-1\right\}$$

$$+ \sum_{k=j_n}^{\infty} q^{k^2/2}(-zq^{-\beta}e^{2\pi i\lambda})^k f(k,n) q^{-ka_n}$$

$$= \sum_{k=-\infty}^{-1} q^{k^2/2}(-q^\beta z^{-1}e^{-2\pi i\lambda})^k + s_{21} + s_{22} + s_{23} + s_{24}.$$

Then,

(180)

$$|s_{21}+s_{24}| \le 2\sum_{k=j_n}^{\infty} q^{k^2/2}(|z|q^{-1-\beta})^k \le 2\,|z|^{j_n}\,q^{j_n^2/2-(\beta+1)j_n}\sum_{k=0}^{\infty} q^{k^2/2}(|z|q^{-1-\beta+j_n})^k$$

$$\le 2\theta_3(|z|^{-1}q^\beta;\sqrt{q})\,|z|^{j_n}\,q^{j_n^2/2-(\beta+1)j_n},$$

$$|s_{22}| \le \frac{2q^{k_n}}{1-q}\sum_{k=1}^{\infty} q^{k^2/2}(|z|q^{-\beta})^k \le \frac{2q^{k_n}}{1-q}\theta_3(|z|^{-1}q^\beta;\sqrt{q}), \tag{181}$$

and

$$|s_{23}| \le \frac{3\log n}{n^\rho}\sum_{k=0}^{\infty} q^{k^2/2}(|z|q^{-\beta})^k \le \frac{3\log n}{n^\rho}\theta_3(|z|^{-1}q^\beta;\sqrt{q}) \tag{182}$$

for $n$ sufficiently large. Hence

$$\frac{(q;q)_\infty s_2}{(-ze^{2\pi i n\theta})^m q^{m(n\tau+m/2)}} = \sum_{k=-\infty}^{-1} q^{k^2/2}(-q^\beta z^{-1}e^{-2\pi i\lambda})^k + r_2(n), \tag{183}$$

where

$$r_2(n) = s_{21} + s_{22} + s_{23} + s_{24}, \tag{184}$$

and

$$|r_2(n)| \leq 3\theta_3(|z|^{-1}q^{\beta}; \sqrt{q}) \left\{ \frac{\log n}{n^{\rho}} + \frac{q^{k_n}}{1-q} + |z|^{j_n}\, q^{j_n^2/2-(\beta+1)j_n} \right\} \tag{185}$$

for $n$ sufficiently large. Therefore,

$$\frac{E_q(-q^{ns+1/2}z)(q;q)_\infty}{(-ze^{2\pi i n\theta})^m q^{m(n\tau+m/2)}} = \theta_4\left(z^{-1}q^{\beta}e^{-2\pi i\lambda}; \sqrt{q}\right) + r_{qe}(n|6), \tag{186}$$

with

(187)

$$|r_{qe}(n|6)| \leq 6\theta_3(|z|^{-1}q^{\beta}; \sqrt{q}) \left\{ \frac{\log n}{n^{\rho}} + \frac{q^{k_n}}{1-q} + |z|^{j_n}\, q^{j_n^2/2-(\beta+1)j_n} + \frac{q^{j_n^2/2+(\beta-1)j_n}}{|z|^{j_n}} \right\}$$

for $n$ sufficiently large.

2.2.6. *Proof for case 7.* Observe that

(188)

$$\begin{aligned}
&\frac{(q;q)_\infty s_1}{(-ze^{2\pi i n\theta})^m q^{m(n\tau+m/2)}} = \sum_{k=0}^{m} q^{k^2/2}(-q^{\beta_1}z^{-1}e^{-2\pi i\beta_2})^k q^{ka_n}e^{-2\pi i k b_n}e(k,n) \\
&= \sum_{k=0}^{\infty} q^{k^2/2}(-q^{\beta_1}z^{-1}e^{-2\pi i\beta_2})^k - \sum_{k=j_n}^{\infty} q^{k^2/2}(-q^{\beta_1}z^{-1}e^{-2\pi i\beta_2})^k \\
&+ \sum_{k=0}^{j_n-1} q^{k^2/2}(-q^{\beta_1}z^{-1}e^{-2\pi i\beta_2})^k \left\{q^{ka_n}-1\right\} + \sum_{k=0}^{j_n-1} q^{k^2/2}(-q^{\beta_1}z^{-1}e^{-2\pi i\beta_2})^k q^{ka_n}\left\{e(k,n)-1\right\} \\
&+ \sum_{k=0}^{j_n-1} q^{k^2/2}(-q^{\beta_1}z^{-1}e^{-2\pi i\beta_2})^k q^{ka_n}\left\{e^{-2\pi i k b_n}-1\right\} e(k,n) \\
&+ \sum_{k=j_n}^{m} q^{k^2/2}(-q^{\beta_1}z^{-1}e^{-2\pi i\beta_2})^k q^{ka_n}e^{-2\pi i k b_n}e(k,n) \\
&= \sum_{k=0}^{\infty} q^{k^2/2}(-q^{\beta_1}z^{-1}e^{-2\pi i\beta_2})^k + s_{11} + s_{12} + s_{13} + s_{14} + s_{15}.
\end{aligned}$$

For $n$ sufficiently large we have

$$\begin{aligned}
|s_{11} + s_{15}| &\leq 2\sum_{k=j_n}^{\infty} q^{k^2/2}(q^{\beta_1-1}|z|^{-1})^k \leq \frac{2q^{j_n^2/2+(\beta_1-1)j_n}}{|z|^{j_n}} \sum_{k=0}^{\infty} q^{k^2/2}(q^{\beta_1-1+j_n}|z|^{-1})^k \\
&\leq \frac{2q^{j_n^2/2+(\beta_1-1)j_n}}{|z|^{j_n}}\theta_3(|z|^{-1}q^{\beta_1}; \sqrt{q}),
\end{aligned} \tag{189}$$

$$|s_{12}| \leq \frac{3\log n}{n^{\rho}} \sum_{k=0}^{\infty} q^{k^2/2}(q^{\beta_1}|z|^{-1})^k \leq \frac{3\log n}{n^{\rho}}\theta_3(|z|^{-1}q^{\beta_1}; \sqrt{q}), \tag{190}$$

$$|s_{13}| \le \frac{2q^{k_n}}{1-q}\sum_{k=0}^{\infty} q^{k^2/2}(q^{\beta_1}|z|^{-1})^k \le \frac{2q^{k_n}}{1-q}\theta_3(|z|^{-1}q^{\beta_1};\sqrt{q}), \tag{191}$$

and

$$|s_{14}| \le \frac{24\log n}{n^{\rho}}\theta_3(|z|^{-1}q^{\beta_1};\sqrt{q}). \tag{192}$$

Hence,

$$\frac{(q;q)_\infty s_1}{(-ze^{2\pi i n\theta})^m q^{m(n\tau+m/2)}} = \sum_{k=0}^{\infty} q^{k^2/2}(-q^{\beta_1}z^{-1}e^{-2\pi i\beta_2})^k + r_1(n), \tag{193}$$

where,

$$r_1(n) = s_{11} + s_{12} + s_{13} + s_{14} + s_{15}, \tag{194}$$

and

$$|r_1(n)| \le 27\theta_3(|z|^{-1}q^{\beta_1};\sqrt{q})\left\{\frac{q^{j_n^2/2+(\beta_1-1)j_n}}{|z|^{j_n}} + \frac{q^{k_n}}{1-q} + \frac{\log n}{n^{\rho}}\right\} \tag{195}$$

for $n$ sufficiently large.

Similarly,

$$\begin{aligned}
&\frac{(q;q)_\infty s_2}{(-ze^{2\pi i n\theta})^m q^{m(n\tau+m/2)}} = \sum_{k=1}^{\infty} q^{k^2/2}(-zq^{-\beta_1}e^{2\pi i\beta_2})^k f(k,n)e^{2\pi i k b_n}q^{-ka_n} \\
&= \sum_{k=-\infty}^{-1} q^{k^2/2}(-q^{\beta_1}z^{-1}e^{-2\pi i\beta_2})^k - \sum_{k=j_n}^{\infty} q^{k^2/2}(-zq^{-\beta_1}e^{2\pi i\beta_2})^k \\
&+ \sum_{k=1}^{j_n-1} q^{k^2/2}(-zq^{-\beta_1}e^{2\pi i\beta_2})^k\left\{f(k,n)-1\right\} + \sum_{k=1}^{j_n-1} q^{k^2/2}(-zq^{-\beta_1}e^{2\pi i\beta_2})^k f(k,n)\left\{q^{-ka_n}-1\right\} \\
&+ \sum_{k=1}^{j_n-1} q^{k^2/2}(-zq^{-\beta_1}e^{2\pi i\beta_2})^k f(k,n)q^{-ka_n}\left\{e^{2\pi i k b_n}-1\right\} \\
&+ \sum_{k=j_n}^{\infty} q^{k^2/2}(-zq^{-\beta_1}e^{2\pi i\beta_2})^k f(k,n)e^{2\pi i k b_n}q^{-ka_n} \\
&= \sum_{k=-\infty}^{-1} q^{k^2/2}(-q^{\beta_1}z^{-1}e^{-2\pi i\beta_2})^k + s_{21} + s_{22} + s_{23} + s_{24} + s_{25}.
\end{aligned} \tag{196}$$

For $n$ sufficiently large we have

$$\begin{aligned}
|s_{21} + s_{25}| &\le 2\sum_{k=j_n}^{\infty} q^{k^2/2}(|z|q^{-\beta_1-1})^k \le 2q^{j_n^2/2-(\beta_1+1)j_n}|z|^{j_n}\sum_{k=0}^{\infty} q^{k^2/2}(|z|q^{j_n-\beta_1-1})^k \\
&\le 2q^{j_n^2/2-(\beta_1+1)j_n}|z|^{j_n}\theta_3(|z|^{-1}q^{\beta_1};\sqrt{q}),
\end{aligned} \tag{197}$$

$$|s_{22}| \le \frac{2q^{k_n}}{1-q}\sum_{k=1}^{\infty} q^{k^2/2}(|z|q^{-\beta_1})^k \le \frac{2q^{k_n}}{1-q}\theta_3(|z|^{-1}q^{\beta_1};\sqrt{q}), \tag{198}$$

$$|s_{23}| \leq \frac{3\log n}{n^\rho}\sum_{k=1}^{\infty} q^{k^2/2}(|z|q^{-\beta_1})^k \leq \frac{3\log n}{n^\rho}\theta_3(|z|^{-1}q^{\beta_1};\sqrt{q}) \tag{199}$$

and

$$|s_{24}| \quad \leq \quad \frac{24\log n}{n^\rho}\theta_3(|z|^{-1}q^{\beta_1};\sqrt{q}). \tag{200}$$

Hence,

$$\frac{(q;q)_\infty s_2}{(-ze^{2\pi in\theta})^m q^{m(n\tau+m/2)}} = \sum_{k=-\infty}^{-1} q^{k^2/2}(-q^{\beta_1}z^{-1}e^{-2\pi i\beta_2})^k + r_2(n), \tag{201}$$

where,

$$r_2(n) = s_{21}+s_{22}+s_{23}+s_{24}+s_{25}, \tag{202}$$

and

$$|r_2(n)| \leq 27\theta_3(|z|^{-1}q^{\beta_1};\sqrt{q})\left\{\frac{\log n}{n^\rho}+\frac{q^{k_n}}{1-q}+q^{j_n^2/2-(\beta_1+1)j_n}|z|^{j_n}\right\}. \tag{203}$$

Therefore,

$$\frac{E_q(-q^{ns+1/2}z)(q;q)_\infty}{(-ze^{2\pi in\theta})^m q^{m(n\tau+m/2)}} = \theta_4\left(z^{-1}q^{\beta_1}e^{-2\pi i\beta_2};\sqrt{q}\right)+r_{qe}(n|7), \tag{204}$$

where,

$$r_{qe}(n|7) = r_1(n)+r_2(n), \tag{205}$$

with

$$|r_{qe}(n|7)| \leq 54\theta_3(|z|^{-1}q^{\beta_1};\sqrt{q})\left\{\frac{\log n}{n^\rho}+\frac{q^{k_n}}{1-q}+q^{j_n^2/2-(\beta_1+1)j_n}|z|^{j_n}+\frac{q^{j_n^2/2+(\beta_1-1)j_n}}{|z|^{j_n}}\right\} \tag{206}$$

for $n$ sufficiently large.

### 2.3. **Proofs For Corollary 2.2.**

2.3.1. *Proof for case 1.* We notice that

$$\frac{q^{n\tau+1/2}}{1-q} = \mathcal{O}\left(n^a e^{-2\pi\tau n^{1-a}}\right) \tag{207}$$

for $n$ sufficiently large. Then from (92) and (93) we have,

$$E_q(-\exp 2\pi(u-n^{1-a}\tau-n^{-a}/2)) = 1+\mathcal{O}\left(n^a e^{-2\pi\tau n^{1-a}}\right) \tag{208}$$

as $n\to\infty$.

2.3.2. *Proof for case 2.* From (91), 10 and (98) we have

$$\left(\frac{q^{k_n}}{1-q}+\frac{q^{k_n^2/2+\lambda k_n}}{|z|^{k_n}}\right)e^{\pi n^a/4}=\mathcal{O}\left(e^{-2\pi n^a}\right) \tag{209}$$

as $n \to \infty$. From (31) and (32) we have

$$\begin{aligned}\theta_3\left(z^{-1}q^{\lambda};\sqrt{q}\right)&=\theta_3(n^{-a}\lambda i+ui|n^{-a}i)\\&=\sqrt{n^a}e^{\pi n^{-a}(n^a u+\lambda)^2}\theta_3(n^a u+\lambda|n^a i)\\&=\sqrt{n^a}e^{\pi n^{-a}(n^a u+\lambda)^2}\left\{1+\mathcal{O}\left(e^{-\pi n^a}\right)\right\},\end{aligned} \tag{210}$$

and

$$\begin{aligned}\theta_4\left(z^{-1}q^{\lambda};\sqrt{q}\right)&=\theta_4(n^{-a}\lambda i+ui|n^{-a}i)\\&=\sqrt{n^a}e^{\pi n^{-a}(n^a u+\lambda)^2}\theta_2(n^a u+\lambda|n^a i)\\&=2\sqrt{n^a}\exp\left(\pi n^{-a}(n^a u+\lambda)^2-\frac{\pi n^a}{4}\right)\\&\cos\pi(n^a u+\lambda)\left\{1+\mathcal{O}\left(e^{-2\pi n^a}\right)\right\},\end{aligned} \tag{211}$$

and

$$\frac{1}{(q;q)_\infty}=\frac{\exp\left(\pi n^a/12-\pi n^{-a}/12\right)}{\sqrt{n^a}}\left\{1+\mathcal{O}\left(e^{-2\pi n^a}\right)\right\} \tag{212}$$

for $n$ sufficiently large.

Then,

$$\begin{aligned}&E_q(-\exp 2\pi(u-n^{1-a}\tau-n^{-a}/2))\\&=\frac{2\exp\left(\pi n^{-a}(n^a u-n\tau)^2\right)\left\{\cos\pi(n^a u+\lambda)+\mathcal{O}\left(e^{-2\pi n^a}\right)\right\}}{(-1)^{\lambda+n\tau}\exp\left(\pi n^a/6+\pi n^{-a}/12\right)}\end{aligned} \tag{213}$$

for $n$ sufficiently large.

### 2.4. **Proof for Corollary 2.3.**

2.4.1. *Proof for case 1.* Since

$$q^{n\tau+1/2}=\mathcal{O}\left(\log n\exp\left(-\frac{2n\pi\tau}{\gamma\log n}\right)\right) \tag{214}$$

as $n \to \infty$, then

$$E_q\left(-\exp 2\pi\left(u-\frac{n\tau+1/2}{\gamma\log n}\right)\right)=1+\mathcal{O}\left(\log n\exp\left(-\frac{2n\pi\tau}{\gamma\log n}\right)\right) \tag{215}$$

for $n$ sufficiently large.

2.4.2. *Proof for case 2.* In this case we have

$$\begin{aligned}\theta_3(z^{-1}q^{\lambda};\sqrt{q}) &= \theta_3\left(ui+\frac{\lambda i}{\gamma\log n}\mid\frac{i}{\gamma\log n}\right)\\ &= \sqrt{\gamma\log n}\exp\frac{(\gamma u\log n+\lambda)^2}{\gamma\log n}\theta_3\left(\gamma u\log n+\lambda\mid i\gamma\log n\right)\\ &= \sqrt{\gamma\log n}\exp\frac{(\gamma u\log n+\lambda)^2}{\gamma\log n}\left\{1+\mathcal{O}(n^{-\pi\gamma})\right\},\end{aligned}\tag{216}$$

and

$$\begin{aligned}\theta_4(z^{-1}q^{\lambda};\sqrt{q}) &= \theta_4\left(ui+\frac{\lambda i}{\gamma\log n}\mid\frac{i}{\gamma\log n}\right)\\ &= \sqrt{\gamma\log n}\exp\frac{(\gamma u\log n+\lambda)^2}{\gamma\log n}\theta_2\left(\gamma u\log n+\lambda\mid i\gamma\log n\right)\\ &= 2n^{-\pi\gamma/4}\sqrt{\gamma\log n}\exp\frac{(\gamma u\log n+\lambda)^2}{\gamma\log n}\\ &\times\cos\pi(\gamma u\log n+\lambda)\left\{1+\mathcal{O}(n^{-2\pi\gamma})\right\},\end{aligned}\tag{217}$$

and

$$\frac{1}{(q;q)_\infty}=\frac{n^{\gamma\pi/12}\exp\left(-\frac{\pi}{12\gamma\log n}\right)}{\sqrt{\gamma\log n}}\left\{1+\mathcal{O}(n^{-2\pi\gamma})\right\}\tag{218}$$

as $n\to\infty$. Thus,

$$\begin{aligned}E_q\left(-\exp 2\pi\left(u-\frac{n\tau+1/2}{\gamma\log n}\right)\right) &= \exp\left(\frac{\pi(\gamma u\log n-n\tau)^2}{\gamma\log n}-\frac{\pi}{12\gamma\log n}\right)\\ &\times\frac{2\left\{\cos\pi(\gamma u\log n+\lambda)+\mathcal{O}(n^{-2\pi\gamma})\right\}}{(-1)^{\tau n+\lambda}n^{\gamma\pi/6}}\end{aligned}\tag{219}$$

as $n\to\infty$.

2.4.3. *Proof for case 3.* A similar calculation to get

$$\begin{aligned}E_q\left(-\exp 2\pi\left(u-\frac{n\tau+1/2}{\gamma\log n}\right)\right) &= \exp\left(\frac{\pi(\gamma u\log n-n\tau)^2}{\gamma\log n}-\frac{\pi}{12\gamma\log n}\right)\\ &\times\frac{2\left\{\cos\pi(\gamma u\log n+\beta)+\mathcal{O}(n^{-8\rho/9}\log n)\right\}}{(-1)^{m}n^{2\rho/27}}\end{aligned}\tag{220}$$

as $n\to\infty$, where

$$\gamma=\frac{4\rho}{9\pi}.\tag{221}$$

## 3. Ramanujan Functions

Ramanujan function $A_q(z)$ (**??**), which is also called $q$-Airy function in the literature, appears repeatedly in Ramanujan's work starting from the Rogers-Ramanujan identities, where $A_q(-1)$ and $A_q(-q)$ are expressed as infinite products, [3], to properties of and conjectures about its zeros, [5, 6, 10, 16]. It is called $q$-Airy function because it appears repeatedly in the Plancherel-Rotach type asymptotics of $q$-orthogonal polynomials, just like the classical Airy function in the classical Plancherel-Rotach asymptotics of classical orthogonal polynomials [25, 13, 7, 8].

Apply inequalities in (13), we have

$$\left|A_q'(z)\right| \le \frac{q}{1-q}A_q(-|z|), \quad z\in\mathbb{C}, \tag{222}$$

and

$$|A_q((1-q)z)| \le e^{q|z|}, \quad z\in\mathbb{C}, \tag{223}$$

or

$$|A_q(z)| \le e^{q|z|/(1-q)}, \quad z\in\mathbb{C}. \tag{224}$$

Similar to $E_q(z)$ case, we could prove that

$$\lim_{q\to 1} A_q((1-q)z) = e^{-z}, \quad z\in\mathbb{C}. \tag{225}$$

Hence, $A_q(z)$ is really one of many $q$-analogues of the exponential function $e^{-z}$.

For any nonzero complex number $z$, then,

$$|A_q(z)| \le \sum_{k=0}^{\infty} \frac{q^{k^2/2}}{(q;q)_k}\left(q^{k/2}\,|z|\right)^k, \tag{226}$$

clearly, the terms $q^{k^2/2}|z|^k$ are bounded by

$$\exp\left\{-\frac{\log^2|z|}{2\log q}\right\}, \tag{227}$$

for $k = 0, 1, ...$, thus,

$$\begin{aligned} |A_q(z)| &\le \exp\left\{-\frac{\log^2|z|}{2\log q}\right\}\sum_{k=0}^{\infty}\frac{q^{k^2/2}}{(q;q)_k} \\ &\le (-\sqrt{q};q)_\infty \exp\left\{-\frac{\log^2|z|}{2\log q}\right\}, \end{aligned} \tag{228}$$

or we have

$$|A_q(z)| \le (-\sqrt{q};q)_\infty \exp\left\{-\frac{\log^2|z|}{2\log q}\right\} \tag{229}$$

for any nonzero complex number $z$.

### 3.1. **Asymptotic Formulas for $A_q(z)$.** Assume that

$$s = 2\tau + \frac{2\pi i\theta}{\log q}, \quad \tau,\theta\in\mathbb{R}, \tag{230}$$

then for any complex number $z$ we have

$$A_q(q^{ns}z) = \sum_{k=0}^{\infty}\frac{q^{k^2}}{(q;q)_k}(-q^{2n\tau}ze^{2n\pi i\theta})^k. \tag{231}$$

**Theorem.** *For any nonzero complex number $z$, let*

$$j_n = \left\lfloor \frac{q^4\log n}{-\log q}\right\rfloor, \quad k_n = \left\lfloor\frac{-n\tau}{2}\right\rfloor, \tag{232}$$

*we have the following for Ramanujan function $A_q(z)$:*

(1) *If* $\tau > 0$, *then,*

$$A_q(q^{ns}z) = 1 + r_{rf}(n|1), \tag{233}$$

*and*

$$|r_{rf}(n|1)| \le \frac{|z|q^{2n\tau+1}}{1-q} \exp\left(\frac{|z|q^{2n\tau+1}}{1-q}\right). \tag{234}$$

(2) *Assume that* $\tau = 0$, *if for some fixed real numbers* $\beta$ *and* $\rho \ge 1$ *there are infinitely many positive integers* $n$ *such that*

$$n\theta = m + \beta + b_n, \quad |b_n| < \frac{1}{n^\rho}, \quad m \in \mathbb{Z}, \tag{235}$$

*then,*

$$A_q(q^{ns}z) = A_q\left(ze^{2\pi i\beta}\right) + r_{rf}(n|3), \tag{236}$$

*and*

$$|r_{rf}(n|3)| \le 24 \exp\left(\frac{|z|q}{1-q}\right)\left\{\frac{\log n}{n^\rho} + \frac{q^{j_n}|z|^{j_n}}{j_n!(1-q)^{j_n}}\right\} \tag{237}$$

*for* $n$ *sufficiently large.*

(3) *Assume that* $\tau < 0$. *If for some fixed rational numbers* $\lambda$ *and* $\lambda_1$ *there are infinitely many positive integers* $n$ *such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}, \quad n\theta = m_1 + \lambda_1, \quad m_1 \in \mathbb{Z}, \tag{238}$$

*then,*

$$\frac{A_q(q^{ns}z)(q;q)_\infty}{(-ze^{2\pi in\theta})^m q^{m(2n\tau+m)}} = \theta_4\left(z^{-1}q^{2\lambda}e^{-2\pi i\lambda_1};q\right) + r_{rf}(n|4), \tag{239}$$

*and*

$$|r_{rf}(n|4)| \le 4\theta_3\left(|z|^{-1}q^{2\lambda};q\right)\left\{\frac{q^{k_n}}{1-q} + \frac{q^{k_n^2+2\lambda k_n}}{|z|^{k_n}}\right\} \tag{240}$$

*for* $n$ *sufficiently large.*

(4) *Assume that* $\tau < 0$. *If for some fixed real numbers* $\beta$, $\lambda$ *and* $\rho \ge 1$ *there exists infinitely many positive integers* $n$ *such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}, \quad n\theta = m_1 + \beta + b_n, \quad |b_n| < \frac{1}{n^\rho}, \quad m_1 \in \mathbb{Z}, \tag{241}$$

*then,*

$$\frac{A_q(q^{ns}z)(q;q)_\infty}{(-ze^{2\pi in\theta})^m q^{m(2n\tau+m)}} = \theta_4\left(z^{-1}q^{2\lambda}e^{-2\pi i\beta};q\right) + r_{rf}(n|5), \tag{242}$$

*and*

$$|r_{rf}(n|5)| \le 48\theta_3\left(|z|^{-1}q^{2\lambda};q\right)\left\{\frac{q^{j_n^2+2\lambda j_n}}{|z|^{j_n}} + |z|^{j_n}q^{j_n^2-2\lambda j_n} + \frac{q^{k_n}}{1-q} + \frac{\log n}{n^\rho}\right\} \tag{243}$$

*for* $n$ *sufficiently large.*

(5) *Assume that* $\tau < 0$. *If for some fixed real numbers* $\lambda$, $\beta$ *and* $\rho \geq 1$ *there exist infinitely many positive integers* $n$ *such that*

$$-n\tau = m + \beta + a_n, \quad |a_n| < \frac{1}{n^\rho}, \quad m \in \mathbb{N}, \quad n\theta = m_1 + \lambda \quad m_1 \in \mathbb{Z}, \tag{244}$$

*then,*

$$\frac{A_q(q^{ns}z)(q;q)_\infty}{(-ze^{2\pi in\theta})^m q^{m(2n\tau+m)}} = \theta_4\left(z^{-1}q^{2\beta}e^{-2\pi i\lambda};q\right) + r_{rf}(n|6), \tag{245}$$

*and*

$$|r_{rf}(n|6)| \leq 12\theta_3\left(|z|^{-1}q^{2\beta};q\right)\left\{\frac{q^{j_n^2+2(\beta-1)j_n}}{|z|^{j_n}} + |z|^{j_n}\,q^{j_n^2-2(\beta+1)j_n} + \frac{q^{k_n}}{1-q} + \frac{\log n}{n^\rho}\right\} \tag{246}$$

*for* $n$ *sufficiently large.*

(6) *Assume that* $\tau < 0$. *If for some fixed real numbers* $\beta_1, \beta_2$ *and* $\rho > 0$ *there exists infinitely many positive integers* $n$ *such that*

$$-n\tau = m + \beta_1 + a_n, \quad m \in \mathbb{N}, \quad |a_n| < \frac{1}{n^\rho}, \tag{247}$$

*and*

$$n\theta = m_1 + \beta_2 + b_n, \quad m_1 \in \mathbb{Z}, \quad |b_n| < \frac{1}{n^\rho}, \tag{248}$$

*then,*

$$\frac{A_q(q^{ns}z)(q;q)_\infty}{(-ze^{2\pi in\theta})^m q^{m(2n\tau+m)}} = \theta_4\left(z^{-1}q^{2\beta_1}e^{-2\pi i\beta_2};q\right) + r_{rf}(n|7), \tag{249}$$

*and*

$$|r_{rf}(n|7)| \leq 54\theta_3\left(|z|^{-1}q^{2\beta_1};q\right)\left\{\frac{q^{j_n^2+2(\beta_1-1)j_n}}{|z|^{j_n}} + |z|^{j_n}\,q^{j_n^2-2(\beta_1+1)j_n} + \frac{q^{k_n}}{1-q} + \frac{\log n}{n^\rho}\right\} \tag{250}$$

*for* $n$ *sufficiently large.*

In the following corollaries, we let

$$z := e^{2\pi u}, \quad u \in \mathbb{R}. \tag{251}$$

**Corollary 3.1.** *Assume that*

$$0 < a < \frac{1}{2}, \quad n \in \mathbb{N}, \quad q = \exp(-\pi n^{-a}), \tag{252}$$

*we have the following results for* $A_q(z)$:

(1) *Assume that* $\tau > 0$, *we have*

$$A_q(\exp 2\pi(u - n^{1-a}\tau)) = 1 + \mathcal{O}\left(n^a \exp(-2\pi\tau n^{1-a})\right) \tag{253}$$

*for* $n$ *sufficiently large.*

(2) *Assume that* $\tau < 0$, *if for some fixed real number* $\lambda$, *there are infinitely many* $m$ *and* $n > 0$ *such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}. \tag{254}$$

*Then,*

$$\begin{aligned} &A_q\left(\exp\left(2\pi(u-n^{1-a}\tau)\right)\right)\\ &=\frac{\sqrt{2}\exp\left\{\pi n^{-a}(n^a u-\tau n)^2\right\}\left\{\cos\pi(n^a u+\lambda)+\mathcal{O}(e^{-2\pi n^a})\right\}}{(-1)^{\tau n+\lambda}\exp\left\{\frac{\pi n^{-a}}{24}+\frac{\pi n^a}{12}\right\}}\end{aligned}\tag{255}$$

*as $n$ sufficiently large.*

Similarly, we have the following:

**Corollary 3.2.** *Assume that*

$$q=\exp(-\frac{\pi}{\gamma\log n}),\quad n\geq 2,\quad \gamma>0,\tag{256}$$

*we have the following results for $A_q(z)$:*

1. *Assume that $\tau>0$, we have*

$$A_q\left(\exp 2\pi\left(u-\frac{n\tau}{\gamma\log n}\right)\right)=1+\mathcal{O}\left(\log n\exp\left(-\frac{2\pi n\tau}{\gamma\log n}\right)\right)\tag{257}$$

   *for $n$ sufficiently large.*

2. *Assume that $\tau<0$ and for some fixed real number $\lambda$, there are infinitely many $m$ and $n>0$ such that*

$$-n\tau=m+\lambda,\quad m\in\mathbb{N}.\tag{258}$$

   *Then,*

$$\begin{aligned} &A_q\left(\exp 2\pi\left(u-\frac{n\tau}{\gamma\log n}\right)\right)\\ &=\sqrt{2}\frac{\exp\left(\frac{\pi(u\gamma\log n-\tau n)^2}{\gamma\log n}\right)\left\{\cos\pi(u\gamma\log n+\lambda)+\mathcal{O}(n^{-2\pi\gamma})\right\}}{(-1)^{n\tau+\lambda}n^{\pi\gamma/12}\exp\left(\frac{\pi}{24\gamma\log n}\right)}\end{aligned}\tag{259}$$

   *for $n$ sufficiently large.*

3. *Assume that $\tau<0$, if for some fixed real numbers $\beta$, $\rho>0$ and $\lambda$ there exist infinitely many positive integers $n$ such that*

$$-n\tau=m+\beta+a_n,\quad |a_n|<\frac{1}{n^\rho},\quad m\in\mathbb{N},\tag{260}$$

   *then for each of such $n$, we have*

$$\begin{aligned} &A_q\left(\exp 2\pi\left(u-\frac{n\tau}{\gamma\log n}\right)\right)\\ &=\sqrt{2}\frac{\exp\left(\frac{\pi(u\gamma\log n-\tau n)^2}{\gamma\log n}\right)\left\{\cos\pi(u\gamma\log n+\beta)+\mathcal{O}(n^{-8\rho/9}\log n)\right\}}{(-1)^m n^{\rho/27}\exp\left(\frac{\pi}{24\gamma\log n}\right)}\end{aligned}\tag{261}$$

   *for $n$ sufficiently large, where*

$$\gamma=\frac{4\rho}{9\pi}.\tag{262}$$

3.1.1. *Proof for Theorem 4.1.* Assume that $\tau < 0$ and

$$-n\tau = m + c_n, \quad m \in \mathbb{N}, \quad n\theta = m_1 + d_n, \quad m_1 \in \mathbb{Z}, \tag{263}$$

then,

$$\begin{aligned} A_q(q^{ns}z) &= \sum_{k=0}^{\infty} \frac{q^{k^2}}{(q;q)_k} \left(-ze^{2\pi in\theta}q^{2n\tau}\right)^k \\ &= \sum_{k=0}^{m} \frac{q^{k^2}}{(q;q)_k} \left(-ze^{2\pi in\theta}q^{2n\tau}\right)^k + \sum_{k=m+1}^{\infty} \frac{q^{k^2}}{(q;q)_k} \left(-ze^{2\pi in\theta}q^{2n\tau}\right)^k \\ &= s_1 + s_2. \end{aligned} \tag{264}$$

Reverse summation order in $s_1$,

$$\frac{(q;q)_\infty s_1}{(-ze^{2\pi in\theta})^m q^{m(2n\tau+m)}} = \sum_{k=0}^{m} q^{k^2}(-q^{2c_n}z^{-1}e^{-2\pi id_n})^k e(k,n), \tag{265}$$

where

$$e(k,n) = \frac{(q;q)_\infty}{(q;q)_{m-k}}, \tag{266}$$

then

$$|e(k,n)| \le 1 \tag{267}$$

for $0 \le k \le m$. From Lemma 1.1 we have

$$|e(k,n) - 1| = |r_1(q; m-k)| \le \frac{2q^{2+k_n}}{1-q}. \tag{268}$$

for $0 \le k \le k_n - 1$ and $n$ sufficiently large.

We shift the summation index from $k$ to $k+m$ in $s_2$,

$$\frac{(q;q)_\infty s_2}{(-ze^{2\pi in\theta})^m q^{m(2n\tau+m)}} = \sum_{k=1}^{\infty} q^{k^2}(-zq^{-2c_n}e^{2\pi id_n})^k f(k,n), \tag{269}$$

where

$$f(k,n) = \frac{(q;q)_\infty}{(q;q)_{m+k}}, \tag{270}$$

thus,

$$|f(k,n)| \le 1, \tag{271}$$

and

$$|f(k,n) - 1| = |r_1(q; m+k)| \le \frac{2q^{2+k_n}}{1-q} \tag{272}$$

for $k \in \mathbb{N}$ and $n$ sufficiently large. The rest are very similar to the corresponding proofs for $E_q(z)$.

3.2. **Proofs for Corollary 3.1.** Notice that

$$\frac{1}{(q;q)_\infty} = \frac{\exp(\pi n^a/6 - \pi n^{-a}/24)}{\sqrt{2n^a}} \left\{1 + \mathcal{O}(e^{-4\pi n^a})\right\}, \tag{273}$$

and

$$\begin{aligned} \theta_4\left(z^{-1}q^{2\lambda};q\right) &= \theta_4(ui + i\lambda n^{-a}|n^{-a}i) \\ &= \sqrt{n^a}\exp\left\{\frac{\pi(n^a u+\lambda)^2}{n^a}\right\}\theta_2\left(n^a u + \lambda \mid n^a i\right) \\ &= 2\sqrt{n^a}\exp\left\{\frac{\pi(n^a u+\lambda)^2}{n^a} - \frac{\pi n^a}{4}\right\} \\ &\times \cos\pi(n^a u+\lambda)\left\{1 + \mathcal{O}(e^{-2\pi n^a})\right\}, \end{aligned} \tag{274}$$

and

$$\begin{aligned} \theta_3\left(|z|^{-1}q^{2\lambda};q\right) &= \theta_3(ui + \lambda i n^{-a}|n^{-a}i) \\ &= \sqrt{n^a}\exp\left\{\frac{\pi(n^a u+\lambda)^2}{n^a}\right\}\theta_3\left(n^a u + \lambda \mid n^a i\right) \\ &= \sqrt{n^a}\exp\left\{\frac{\pi(n^a u+\lambda)^2}{n^a}\right\}\left\{1 + \mathcal{O}(e^{-\pi n^a})\right\}, \end{aligned} \tag{275}$$

The rest of the proof for this corollary are very similar to the $E_q(z)$ case.

3.3. **Proof for Corollary 3.2.** Observe that

$$\frac{1}{(q;q)_\infty} = \frac{n^{\pi\gamma/6}\exp\left(-\frac{\pi}{24\gamma\log n}\right)}{\sqrt{2\gamma\log n}}\left\{1 + \mathcal{O}(n^{-4\pi\gamma})\right\}, \tag{276}$$

and

$$\begin{aligned} \theta_3(z^{-1}q^{2\lambda};q) &= \theta_3\left(ui + i\frac{\lambda}{\gamma\log n} \mid \frac{i}{\gamma\log n}\right) \\ &= \sqrt{\gamma\log n}\exp\left(\frac{\pi(u\gamma\log n+\lambda)^2}{\gamma\log n}\right)\theta_3(u\gamma\log n + \lambda \mid i\gamma\log n) \\ &= \sqrt{\gamma\log n}\exp\left(\frac{\pi(u\gamma\log n+\lambda)^2}{\gamma\log n}\right)\left\{1 + \mathcal{O}(n^{-\pi\gamma})\right\}, \end{aligned} \tag{277}$$

and

$$\begin{aligned} \theta_4(z^{-1}q^{2\lambda};q) &= \theta_4\left(ui + i\frac{\lambda}{\gamma\log n} \mid \frac{i}{\gamma\log n}\right) \\ &= \sqrt{\gamma\log n}\exp\left(\frac{\pi(u\gamma\log n+\lambda)^2}{\gamma\log n}\right)\theta_2(u\gamma\log n + \lambda \mid i\gamma\log n) \\ &= 2n^{-\pi\gamma/4}\sqrt{\gamma\log n}\exp\left(\frac{\pi(u\gamma\log n+\lambda)^2}{\gamma\log n}\right) \\ &\times \cos\pi(u\gamma\log n+\lambda)\left\{1 + \mathcal{O}(n^{-2\pi\gamma})\right\} \end{aligned} \tag{278}$$

as $n \to \infty$. The rest of proof is similar to the proof for $E_q(z)$ case.

## 4. $q$-Bessel Function of Second Kind

Jackson's $q$-Bessel function of second kind $J_\nu^{(2)}(z;q)$ is defined as [22, 13, 3, 9, 20]

$$J_\nu^{(2)}(z;q) := \frac{(q^{\nu+1};q)_\infty}{(q;q)_\infty}\sum_{k=0}^\infty \frac{q^{k^2+k\nu}(-1)^k}{(q,q^{\nu+1};q)_k}\left(\frac{z}{2}\right)^{2k+\nu}. \tag{279}$$

In this chapter, we always assume that $\nu > -1$. Clearly,

$$\left|J_\nu^{(2)}(z;q)\right| \le \frac{(|z|/2)^\nu}{(q;q)_\infty} A_q\left(-\frac{|z|^2 q^\nu}{4}\right), \tag{280}$$

and

$$\left|J_\nu^{(2)}(z;q)\right| \le \frac{(|z|/2)^\nu}{(q;q)_\infty}\exp\left\{\frac{q^{1+\nu}|z|^2}{4(1-q)}\right\}. \tag{281}$$

for any complex number $z$.

For any nonzero complex number $z$, (223) implies

$$\left|J_\nu^{(2)}(z;q)\right| \le \frac{(-\sqrt{q};q)_\infty}{(q;q)_\infty}\left(\frac{|z|}{2}\right)^\nu \exp\left\{-\frac{\log^2\left(|z|^2 q^\nu/4\right)}{2\log q}\right\}. \tag{282}$$

### 4.1. Asymptotics for $J_\nu^{(2)}(z;q)$ .

From (267) we have

$$\frac{J_\nu^{(2)}(2q^{ns-\nu/2}z;q)(q;q)_\infty}{(q^{\nu+1};q)_\infty\left(q^{ns-\nu/2}z\right)^\nu} = \sum_{k=0}^\infty \frac{q^{k^2}\left(-z^2q^{2ns}\right)^k}{(q,q^{\nu+1};q)_k}. \tag{283}$$

Let

$$s = \tau + \frac{\pi i\theta}{\log q}, \quad \tau,\theta\in\mathbb{R}, \tag{284}$$

then for any complex number $z$ we have

$$\frac{J_\nu^{(2)}(2q^{ns-\nu/2}z;q)(q;q)_\infty}{(q^{\nu+1};q)_\infty\left(q^{n\tau-\nu/2}ze^{n\pi i\theta}\right)^\nu} = \sum_{k=0}^\infty \frac{q^{k^2}(-q^{2n\tau}z^2e^{2n\pi i\theta})^k}{(q,q^{\nu+1};q)_k}. \tag{285}$$

**Theorem.** *For any nonzero complex number $z$, let*

$$j_n = \left\lfloor \frac{q^4\log n}{-\log q}\right\rfloor, \quad k_n = \left\lfloor\frac{-n\tau}{2}\right\rfloor, \tag{286}$$

*we have the following for the function $J_\nu^{(2)}(z;q)$:*

(1) *When $\tau > 0$ we have*

$$\frac{J_\nu^{(2)}(2q^{ns-\nu/2}z;q)(q;q)_\infty}{(q^{\nu+1};q)_\infty\left(q^{ns-\nu/2}z\right)^\nu} = 1 + r_{qb}(n|1), \tag{287}$$

*and*

$$|r_{qb}(n|1)| \le \frac{|z|^2\exp\left(|z|^2q^{2n\tau+1}/(1-q)\right)}{(1-q)(q^{\nu+1};q)_\infty}q^{2n\tau+1}. \tag{288}$$

(2) *Assume that* $\tau = 0$. *If for some fixed real numbers* $\beta$ *and* $\rho \geq 1$ *there are infinitely many positive integers* $n$ *such that*

$$n\theta = m + \beta + b_n, \quad |b_n| < \frac{1}{n^{\rho}}, \quad m \in \mathbb{Z}, \tag{289}$$

*then*

$$\frac{J_{\nu}^{(2)}(2q^{ns-\nu/2}z;q)(q;q)_{\infty}}{\left(q^{n\tau-\nu/2}ze^{n\pi i\theta}\right)^{\nu}} = \frac{J_{\nu}^{(2)}(2q^{-\nu/2}ze^{\pi i(m+\beta)};q)(q;q)_{\infty}}{\left(q^{-\nu/2}ze^{\pi i(m+\beta)}\right)^{\nu}} + r_{qb}(n|3), \tag{290}$$

*and*

$$r_{qb}(n|3)| \leq 24\exp(q|z|^2/(1-q))\left\{\frac{\log n}{n^{\rho}} + \frac{1}{j_n!}\left(\frac{q|z|^2}{1-q}\right)^{j_n}\right\} \tag{291}$$

*for* $n$ *sufficiently large.*

(3) *Assume that* $\tau < 0$. *If for some fixed real numbers* $\lambda$ *and* $\lambda_1$ *there are infinitely many positive integers* $n$ *such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}, \quad n\theta = m_1 + \lambda_1, \quad m_1 \in \mathbb{Z}, \tag{292}$$

*then,*

$$\frac{J_{\nu}^{(2)}(2q^{ns-\nu/2}z;q)(q;q)_{\infty}^{2}e^{-n\pi i\theta(2m+\nu)}}{(-1)^{m}z^{\nu+2m}q^{m(2n\tau+m)+n\nu\tau-\nu^2/2}} = \theta_4\left(z^{-2}q^{2\lambda}e^{-2\pi i\lambda_1};q\right) + r_{qb}(n|4), \tag{293}$$

*and*

$$|r_{qb}(n|4)| \leq 12\theta_3\left(|z|^{-2}q^{2\lambda};q\right)\left\{\frac{q^{k_n}}{(1-q)} + \frac{q^{k_n^2+2\lambda k_n}}{|z|^{2k_n}}\right\} \tag{294}$$

*for* $n$ *sufficiently large.*

(4) *Assume that* $\tau < 0$. *If for some fixed real numbers* $\beta$, $\lambda$ *and* $\rho \geq 1$ *there exist infinitely many positive integers* $n$ *such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}, \quad n\theta = m_1 + \beta + b_n, \quad |b_n| < \frac{1}{n^{\rho}}, \quad m_1 \in \mathbb{Z}, \tag{295}$$

*then,*

$$\frac{J_{\nu}^{(2)}(2q^{ns-\nu/2}z;q)(q;q)_{\infty}^{2}e^{-n\pi i\theta(2m+\nu)}}{(-1)^{m}z^{\nu+2m}q^{m(2n\tau+m)+n\nu\tau-\nu^2/2}} = \theta_4\left(z^{-2}q^{2\lambda}e^{-2\pi i\beta};q\right) + r_{rf}(n|5), \tag{296}$$

*and*

$$|r_{rf}(n|5)| \leq 48\theta_3\left(|z|^{-2}q^{2\lambda};q\right)\left\{\frac{q^{j_n^2+2\lambda j_n}}{|z|^{2j_n}} + |z|^{2j_n}\,q^{j_n^2-2\lambda j_n} + \frac{q^{k_n}}{(1-q)} + \frac{\log n}{n^{\rho}}\right\} \tag{297}$$

*for* $n$ *sufficiently large.*

(5) *Assume that* $\tau < 0$. *If for some fixed real numbers* $\beta$, $\lambda$ *and* $\rho \geq 1$ *there exist infinitely many positive integers* $n$ *such that*

$$-n\tau = m + \beta + a_n, \quad m \in \mathbb{N}, \quad |a_n| < \frac{1}{n^{\rho}}, \quad n\theta = m_1 + \lambda \quad m_1 \in \mathbb{Z}, \tag{298}$$

*then,*

$$\frac{J_{\nu}^{(2)}(2q^{ns-\nu/2}z;q)(q;q)_{\infty}^{2}e^{-n\pi i\theta(2m+\nu)}}{(-1)^{m}z^{\nu+2m}q^{m(2n\tau+m)+n\nu\tau-\nu^2/2}} = \theta_4(q^{2\beta}z^{-2}e^{-2\pi i\lambda};q) + r_{qb}(n|6), \tag{299}$$

*and*

(300)
$$|r_{qb}(n|6)| \leq 12\theta_3(|z|^{-2}q^{2\beta};q)\left\{\frac{q^{j_n^2+2(\beta-1)j_n}}{|z|^{2j_n}} + |z|^{2j_n}q^{j_n^2-2(\beta+1)j_n} + \frac{q^{k_n}}{1-q} + \frac{\log n}{n^\rho}\right\}$$

*for $n$ sufficiently large.*

(6) *Assume that $\tau < 0$. If for some fixed real numbers $\beta_1$, $\beta_2$ and $\rho > 0$ there exist infinitely many positive integers $n$ such that*

$$-n\tau = m + \beta + a_n, \quad m \in \mathbb{N}, \quad |a_n| < \frac{1}{n^\rho}, \tag{301}$$

*and*

$$n\theta = m_1 + \beta + b_n, \quad m_1 \in \mathbb{Z}, \quad |b_n| < \frac{1}{n^\rho}. \tag{302}$$

*Then,*

$$\frac{J_\nu^{(2)}(2q^{ns-\nu/2}z;q)(q;q)_\infty^2 e^{-n\pi i\theta(2m+\nu)}}{(-1)^m z^{\nu+2m}q^{m(2n\tau+m)+n\nu\tau-\nu^2/2}} = \theta_4(q^{2\beta_1}z^{-2}e^{-2\pi i\beta_2};q) + r_{qb}(n|7), \tag{303}$$

*and*

(304)
$$|r_{qb}(n|7)| \leq 156\theta_3(q^{2\beta_1}|z|^{-2};q)\left\{\frac{q^{j_n^2+2(\beta_1-1)j_n}}{|z|^{2j_n}} + |z|^{2j_n}q^{j_n^2-2(\beta_1+1)j_n} + \frac{q^{k_n}}{1-q} + \frac{\log n}{n^\rho}\right\}$$

*for $n$ sufficiently large.*

In the following corollaries we let

$$z := e^{\pi u}, \quad u \in \mathbb{R}. \tag{305}$$

**Corollary 4.1.** *Assume that*

$$q = \exp(-n^{-a}\pi), \quad 0 < a < \frac{1}{2}, \quad n \in \mathbb{N}, \tag{306}$$

*we have the following results for $J_\nu^{(2)}(z;q)$:*

(1) *Assume that $\tau > 0$, we have*

$$\begin{aligned} &J_\nu^{(2)}(2\exp\pi(u-\tau n^{1-a}+\nu n^{-a}/2);\exp(-n^{-a}\pi)) \\ &= \frac{n^{a\nu}\exp(\pi\nu u - \pi\tau n^{1-a} + \nu^2 n^{-a}\pi/2)}{(2\pi)^\nu}\left\{\frac{1}{\Gamma(\nu+1)} + o(1)\right\}. \end{aligned} \tag{307}$$

*for $n$ sufficiently large.*

(2) *Assume that $\tau < 0$ and for some fixed real number $\lambda$ there are infinitely many positive integers $n$ such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}. \tag{308}$$

*Then*

$$\begin{aligned} &J_\nu^{(2)}(2\exp\pi(u-\tau n^{1-a}+\nu n^{-a}/2);\exp(-n^{-a}\pi)) \\ &= \exp\left(\frac{\pi}{n^a}(n^a u + \nu/2 - \tau n)^2 + \frac{\pi n^a}{12} + \frac{\pi\nu^2 n^{-a}}{4} - \frac{\pi n^{-a}}{12}\right) \\ &\times (-1)^{\lambda+n\tau}n^{-a}\left\{\cos\pi(n^a u + \lambda) + \mathcal{O}(e^{-2\pi n^a})\right\} \end{aligned} \tag{309}$$

*for $n$ sufficiently large.*

Similarly, we have the following:

**Corollary 4.2.** *Assume that*

$$q = \exp(-\frac{\pi}{\gamma \log n}), \quad n \geq 2, \quad \gamma > 0, \tag{310}$$

*we have the following results for* $J_\nu^{(2)}(z;q)$*:*

(1) *Assume that* $\tau > 0$*, we have*

$$\begin{aligned} &J_\nu^{(2)}\left(2\exp\pi\left(u - \frac{n\tau}{\gamma\log n} + \frac{\nu}{2\gamma\log n}\right); \exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \\ &= \left(\frac{\pi}{\gamma\log n}\right)^\nu \exp\left(\pi\nu u + \frac{\pi\nu^2}{2\gamma\log n} - \frac{\pi n\nu\tau}{\gamma\log n}\right)\left\{\frac{1}{\Gamma(\nu+1)} + o(1)\right\} \end{aligned} \tag{311}$$

*for $n$ sufficiently large.*

(2) *Assume that* $\tau < 0$ *and for some fixed real number* $\lambda$ *there are infinitely many positive integers $n$ such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}, \tag{312}$$

*then,*

$$\begin{aligned} &J_\nu^{(2)}\left(2\exp\pi\left(u - \frac{n\tau}{\gamma\log n} + \frac{\nu}{2\gamma\log n}\right); \exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \\ &= \exp\left\{\frac{\pi(u\gamma\log n - n\tau + \nu/2)^2}{\gamma\log n} - \frac{\pi}{12\gamma\log n} + \frac{\nu^2\pi}{4\gamma\log n}\right\} \\ &\times \frac{(-1)^{n\tau+\lambda} n^{\pi\gamma/12}}{\sqrt{\gamma\log n}}\left\{\cos\pi(\gamma u\log n + \lambda) + \mathcal{O}(n^{-2\pi\gamma})\right\}, \end{aligned} \tag{313}$$

*for $n$ sufficiently large.*

(3) *Assume that* $\tau < 0$*, if for some fixed real numbers* $\beta$*,* $\rho \geq 1$ *and* $\lambda$ *there exist infinitely many positive integers $n$ such that*

$$-n\tau = m + \beta + a_n, \quad |a_n| < \frac{1}{n^\rho}, \quad m \in \mathbb{N}, \tag{314}$$

*then,*

$$\begin{aligned} &J_\nu^{(2)}\left(2\exp\pi\left(u - \frac{n\tau}{\gamma\log n} + \frac{\nu}{2\gamma\log n}\right); \exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \\ &= \exp\left\{\frac{\pi(u\gamma\log n - n\tau + \nu/2)^2}{\gamma\log n} - \frac{\pi}{12\gamma\log n} + \frac{\nu^2\pi}{4\gamma\log n}\right\} \\ &\times \frac{(-1)^m n^{\rho/27}}{\sqrt{\gamma\log n}}\left\{\cos\pi(\gamma u\log n + \lambda) + \mathcal{O}(n^{-8\rho/9})\right\}, \end{aligned} \tag{315}$$

*for $n$ sufficiently large, where*

$$\gamma = \frac{4\rho}{9\pi}. \tag{316}$$

4.2. **Proofs for Theorem 4.1.** Assume that $\tau < 0$ and

$$-n\tau = m + c_n, \quad m \in \mathbb{N}, \quad n\theta = m_1 + d_n, \quad m_1 \in \mathbb{Z}, \tag{317}$$

then,

$$\begin{aligned}\frac{J_\nu^{(2)}(2q^{ns-\nu/2}z;q)(q;q)_\infty}{(q^{\nu+1};q)_\infty\left(q^{n\tau-\nu/2}ze^{n\pi i\theta}\right)^\nu} &= \sum_{k=0}^{\infty}\frac{q^{k^2}(-q^{2n\tau}z^2e^{2n\pi i\theta})^k}{(q,q^{\nu+1};q)_k}\\ &= \sum_{k=0}^{m}\frac{q^{k^2}(-q^{2n\tau}z^2e^{2n\pi i\theta})^k}{(q,q^{\nu+1};q)_k} + \sum_{k=m+1}^{\infty}\frac{q^{k^2}(-q^{2n\tau}z^2e^{2n\pi i\theta})^k}{(q,q^{\nu+1};q)_k}\\ &= s_1 + s_2,\end{aligned} \tag{318}$$

and reverse summation order in $s_1$,

$$\frac{(q,q^{\nu+1};q)_\infty s_1}{(-z^2e^{2\pi in\theta})^m q^{m(2n\tau+m)}} = \sum_{k=0}^{m} q^{k^2}(-q^{2c_n}z^{-2}e^{-2\pi id_n})^k e(k,n) \tag{319}$$

with

$$e(k,n) = (q^{m-k+1}, q^{\nu+1+m-k};q)_\infty, \tag{320}$$

then

$$|e(k,n)| \le 1 \tag{321}$$

for $0 \le k \le m$. We also have

$$|e(k,n) - 1| \le \frac{6q^{k_n}}{1-q}, \tag{322}$$

for $0 \le k \le k_n - 1$ and $n$ sufficiently large. This could be seen by expanding

$$e(k,n) - 1 = \left\{r_1\left(q;m-k\right)+1\right\}\left\{r_1\left(q^{\nu+1};m-k\right)+1\right\} - 1 \tag{323}$$

and estimating each terms by Lemma (1.1) to obtain

We shift the summation index from $k$ to $k+m$ in $s_2$,

$$\frac{(q,q^{\nu+1};q)_\infty s_2}{(-z^2e^{2\pi in\theta})^m q^{m(2n\tau+m)}} = \sum_{k=1}^{\infty} q^{k^2}(-z^2q^{-2c_n}e^{2\pi id_n})^k f(k,n), \tag{324}$$

and

$$f(k,n) = (q^{m+k+1}, q^{\nu+1+m+k};q)_\infty. \tag{325}$$

Thus,

$$|f(k,n)| \le 1 \tag{326}$$

for $k \in \mathbb{N}$. Expand

$$f(k,n) - 1 = \left\{r_1\left(q;m+k\right)+1\right\}\left\{r_1\left(q^{\nu+1};m+k\right)+1\right\} - 1, \tag{327}$$

and estimate each term by Lemma (1.1) to obtain

$$|f(k,n) - 1| \le \frac{6q^{k_n}}{1-q} \tag{328}$$

for $k \in \mathbb{N}$ and $n$ sufficiently large. The rest of proof is similar to the corresponding proof for $E_q(z)$.

4.3. **Proofs for Corollary 4.2.** The $q$-Gamma function is defined as [6, 9, 13, 20]

$$\Gamma_q(x)=\frac{(q;q)_\infty}{(q^x;q)_\infty}(1-q)^{1-x}\quad x\in\mathbb{C}. \tag{329}$$

It is a $q$-analogue of $\Gamma(x)$,

$$\lim_{q\to1}\Gamma_q(x)=\Gamma(x). \tag{330}$$

Thus

$$\frac{(q^{\nu+1};q)_\infty(1-q)^\nu}{(q;q)_\infty}=\frac{1}{\Gamma(\nu+1)}+o(1) \tag{331}$$

as $n$ sufficiently large.

We also have

$$\begin{aligned}\theta_4(z^{-2}q^{2\lambda};q)&=\theta_4(ui+\lambda n^{-a}i|n^{-a}i)\\&=\sqrt{n^a}\exp\frac{\pi}{n^a}(n^au+\lambda)^2\theta_2\left(n^au+\lambda\mid n^ai\right)\\&=2\sqrt{n^a}\exp\left(\frac{\pi}{n^a}(n^au+\lambda)^2-\frac{\pi n^a}{4}\right)\\&\times\cos\pi(n^au+\lambda)\left\{1+\mathcal{O}(e^{-2\pi n^a})\right\},\end{aligned} \tag{332}$$

and

$$\begin{aligned}\theta_3\left(|z|^{-2}q^{2\lambda};q\right)&=\theta_3(ui+\lambda n^{-a}i|n^{-a}i)\\&=\sqrt{n^a}\exp\frac{\pi}{n^a}(n^au+\lambda)^2\theta_3\left(n^au+\lambda\mid n^ai\right)\\&=\sqrt{n^a}\exp\frac{\pi}{n^a}(n^au+\lambda)^2\left\{1+\mathcal{O}(e^{-\pi n^a})\right\}\end{aligned} \tag{333}$$

as $n\to\infty$. The rest of the proof is similar to corresponding proof for $E_q(z)$ case.

4.4. **Proof for Corollary 4.2.** Notice that

$$\frac{1}{(q;q)_\infty}=\frac{n^{\pi\gamma/6}\left\{1+\mathcal{O}\left(n^{-4\pi\gamma}\right)\right\}}{\sqrt{2\gamma\log n}\exp\left(\frac{\pi}{24\gamma\log n}\right)}, \tag{334}$$

and

$$\begin{aligned}\theta_4(z^{-2}q^{2\lambda};q)&=\theta_4\left(ui+\frac{i\lambda}{\gamma\log n}\mid\frac{i}{\gamma\log n}\right)\\&=\sqrt{\gamma\log n}\exp\frac{\pi(u\gamma\log n+\lambda)^2}{\gamma\log n}\theta_2\left(\gamma u\log n+\lambda\mid i\gamma\log n\right)\\&=2n^{-\pi\gamma/4}\sqrt{\gamma\log n}\exp\left(\frac{\pi(u\gamma\log n+\lambda)^2}{\gamma\log n}\right)\\&\times\cos\pi(\gamma u\log n+\lambda)\left\{1+\mathcal{O}(n^{-2\pi\gamma})\right\},\end{aligned} \tag{335}$$

and

$$\begin{aligned}\theta_3(z^{-2}q^{2\lambda};q) &= \theta_3\left(ui+\frac{i\lambda}{\gamma\log n}\mid\frac{i}{\gamma\log n}\right)\\ &=\sqrt{\gamma\log n}\exp\frac{\pi(u\gamma\log n+\lambda)^2}{\gamma\log n}\theta_3\left(\gamma u\log n+\lambda\mid i\gamma\log n\right)\\ &=\sqrt{\gamma\log n}\exp\frac{\pi(u\gamma\log n+\lambda)^2}{\gamma\log n}\left\{1+\mathcal{O}(n^{-\pi\gamma})\right\}\end{aligned}\tag{336}$$

as $n\to\infty$, and the rest of the proof is similar to the corresponding proof for $E_q(z)$.

## 5. Ismail-Masson Orthogonal Polynomials

Ismail-Masson polynomials $\{h_n(x|q)\}_{n=0}^{\infty}$ are defined as [13]

$$h_n(\sinh\xi|q)=\sum_{k=0}^{n}\frac{(q;q)_nq^{k(k-n)}(-1)^ke^{(n-2k)\xi}}{(q;q)_k(q;q)_{n-k}}.\tag{337}$$

Ismail-Masson polynomials come from an indeterminate moment problem. Ismail-Masson orthogonal polynomials satisfy the following orthogonality

$$\int_{-\infty}^{\infty}h_m(x|q)h_n(x|q)w_{im}(x)dx \quad = \quad q^{-n(n+1)/2}(q;q)_n\delta_{m,n},\tag{338}$$

where

$$w_{im}(x):=q^{1/8}\sqrt{\frac{-2}{\pi\log q}}\exp\left(\frac{2\log^2(x+\sqrt{x^2+1})}{\log q}\right).\tag{339}$$

The corresponding orthonormal Ismail-Masson functions are given by

$$h_n(x\|q):=q^{n(n+1)/4}\sqrt{\frac{w_{im}(x)}{(q;q)_n}}h_n(x|q).\tag{340}$$

Let

$$s=\frac{1+2\tau}{2}+i\frac{\theta\pi}{\log q},\quad \sinh\xi_n:=\frac{q^{-ns}z-q^{ns}z^{-1}}{2},\quad \tau,\theta\in\mathbb{R},\tag{341}$$

for any nonzero complex number $z$, then

$$\frac{h_n(\sinh\xi_n|q)}{z^nq^{-n^2s}}=\sum_{k=0}^{n}\frac{q^{k^2}(q;q)_ne^{2nk\theta\pi i}}{(q;q)_k(q;q)_{n-k}}\left(-\frac{q^{2\tau n}}{z^2}\right)^k.\tag{342}$$

Obviously,

$$|h_n(\sinh\xi_n|q)| \quad\le\quad \frac{|z|^n}{q^{n^2(\tau+1/2)}}\sum_{k=0}^{\infty}\frac{q^{k^2}}{(q;q)_k}\left(\frac{q^{2\tau n}}{|z|^2}\right)^k\le\frac{|z|^nA_q\left(-\frac{q^{2\tau n}}{|z|^2}\right)}{q^{n^2(\tau+1/2)}},\tag{343}$$

*or*

$$|h_n(\sinh\xi_n|q)|\le\frac{(-\sqrt{q};q)_\infty|z|^{(4\tau+1)n}}{q^{2n^2(\tau+1/2)^2}}\exp\left(-\frac{2\log^2|z|}{\log q}\right).\tag{344}$$

### 5.1. Asymptotic Formulas For Ismail-Masson Polynomials.

**Theorem 5.1.** *Given any nonzero complex number $z$, let $s$ and $\xi_n$ be defined as in* (340) *and*

$$j_n = \left\lfloor \frac{q^4 \log n}{\log q^{-1}} \right\rfloor, \quad k_n = \min\left\{ \left\lfloor \frac{(\tau+1)n}{2} \right\rfloor, \left\lfloor \frac{-\tau n}{2} \right\rfloor \right\}, \tag{345}$$

*we have the following results for Ismail-Masson polynomials:*

(1) *Assume that $\tau > 0$, we have*

$$\frac{h_n(\sinh \xi_n|q)}{z^n q^{-n^2 s}} = 1 + r_{im}(n|1), \tag{346}$$

*and*

$$|r_{im}(n|1)| \le \frac{\exp\left(q^{2\tau n+1}/(|z|^2(1-q))\right)}{|z|^2(1-q)} q^{2\tau n+1}. \tag{347}$$

(2) *Assume that $\tau = 0$. If for any fixed real number $\lambda$ there are infinitely many positive integers $n$ such that*

$$n\theta = m + \lambda, \quad m \in \mathbb{Z}, \tag{348}$$

*then,*

$$\frac{h_n(\sinh \xi_n|q)}{z^n q^{-n^2 s}} = A_q\left(\frac{e^{2\lambda\pi i}}{z^2}\right) + r_{im}(n|2), \tag{349}$$

*and*

$$|r_{im}(n|2)| \le 6\frac{A_q(-|z|^{-2})}{(q;q)_\infty}\left\{ q^{n/2} + \frac{q^{\lfloor n/2 \rfloor^2}}{|z|^{2\lfloor n/2 \rfloor}} \right\}. \tag{350}$$

*for $n$ is sufficiently large.*

(3) *Assume that $\tau = 0$. If for fixed real numbers $\beta$ and $\rho \ge 1$ there are infinitely positive integers $n$ such that*

$$n\theta = m + \beta + b_n \quad |b_n| < \frac{1}{n^\rho}, \quad m \in \mathbb{Z}, \tag{351}$$

*then,*

$$\frac{h_n(\sinh \xi_n|q)}{z^n q^{-n^2 s}} = A_q\left(\frac{e^{2\pi i\beta}}{z^2}\right) + e_{im}(n|3), \tag{352}$$

*and*

$$|e_{im}(n|3)| \le 24 \exp\left(\frac{q}{|z|^2(1-q)}\right)\left\{ \frac{\log n}{n^\rho} + \frac{q^{n/2}}{1-q} + \frac{(|z|^2(q^{-1}-1))^{-j_n}}{j_n!} \right\}, \tag{353}$$

*for $n$ sufficiently large.*

(4) *Assume that $-\frac{1}{2} < \tau < 0$. If for some fixed real numbers $\lambda$ and $\lambda_1$ there are infinite number of positive integers $n$ such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}, \quad n\theta = m_1 + \lambda_1, \quad m_1 \in \mathbb{Z}, \tag{354}$$

*then,*

$$h_n(\sinh \xi_n|q) = \frac{z^n q^{-n^2 s + m(2\tau n + m)}}{(q;q)_\infty (-z^2 e^{-2\pi i\lambda_1})^m}\left\{ \theta_4\left(z^2 q^{2\lambda} e^{-2\pi i\lambda_1}; q\right) + r_{im}(n|4) \right\}, \tag{355}$$

*and*

$$|r_{im}(n|4)| \le 28\theta_3\left(|z|^2\, q^{2\lambda}; q\right)\left\{\frac{q^{k_n}}{1-q} + |z|^{2k_n} q^{k_n^2+2\lambda k_n} + |z|^{-2k_n} q^{k_n^2-2\lambda k_n}\right\}. \tag{356}$$

*for $n$ sufficiently large.*

(5) *Assume that $-\frac{1}{2} < \tau < 0$. If for some fixed real numbers $\beta$, $\lambda$ and $\rho \ge 1$ there are infinitely many positive integers $n$ such that*

$$n\theta = m_1 + \beta + b_n, \quad |b_n| < \frac{1}{n^\rho}, \quad m_1 \in \mathbb{Z}, \quad -n\tau = m + \lambda, \quad m \in \mathbb{N}, \tag{357}$$

*then,*

$$h_n(\sinh \xi_n|q) = \frac{z^n q^{-n^2 s + m(2\tau n + m)}}{(-z^2 e^{-2n\theta\pi i})^m (q;q)_\infty}\left\{\theta_4\left(z^2 q^{2\lambda} e^{-2\beta\pi i}; q\right) + r_{im}(n|5)\right\}, \tag{358}$$

*and*

$$|r_{im}(n|5)| \le 24\theta_3(|z|^2 q^{2\lambda}; q)\left\{|z|^{-2j_n} q^{j_n^2-2\lambda j_n} + |z|^{2j_n} q^{j_n^2+2\lambda j_n} + \frac{q^{k_n}}{1-q} + \frac{\log n}{n^\rho}\right\} \tag{359}$$

*for $n$ sufficiently large.*

(6) *Assume that $-\frac{1}{2} < \tau < 0$. If for fixed real numbers $\beta$, $\lambda$ and $\rho \ge 1$ there are infinitely many positive integers $n$ such that*

$$-n\tau = m + \beta + a_n, \quad |a_n| < \frac{1}{n^\rho}, \quad m \in \mathbb{N}, \quad n\theta = m_1 + \lambda, \quad m_1 \in \mathbb{Z}, \tag{360}$$

*then,*

$$h_n(\sinh \xi_n|q) = \frac{z^n q^{-n^2 s + m(2\tau n + m)}}{(-z^2 e^{-2n\theta\pi i})^m (q;q)_\infty}\left\{\theta_4\left(z^2 q^{2\beta} e^{-2\lambda\pi i}; q\right) + r_{im}(n|6)\right\}, \tag{361}$$

*and*

$$|r_{im}(n|6)| \le 12\theta_3(|z|^2 q^{2\beta}; q)\left\{\frac{\log n}{n^\rho} + \frac{q^{k_n}}{1-q} + |z|^{2j_n} q^{j_n^2+2(\beta-1)j_n} + |z|^{-2j_n} q^{j_n^2-2(\beta+1)j_n}\right\} \tag{362}$$

*for $n$ sufficiently large.*

(7) *Assume that $-\frac{1}{2} < \tau < 0$. If for fixed real numbers $\beta_1$, $\beta_2$ and $\rho > 0$ there are infinitely many positive integers $n$ such that*

$$-\tau n = m + \beta_1 + a_n, \quad |a_n| < \frac{1}{n^\rho}, \quad m \in \mathbb{N}, \tag{363}$$

*and*

$$n\theta = m_1 + \beta_2 + b_n, \quad |b_n| < \frac{1}{n^\rho}, \quad m_1 \in \mathbb{Z}. \tag{364}$$

*Then,*

$$h_n(\sinh \xi_n|q) = \frac{z^n q^{-n^2 s + m(2\tau n + m)}}{(-z^2 e^{-2n\theta\pi i})^m (q;q)_\infty}\left\{\theta_4\left(z^2 q^{2\beta_1} e^{-2\beta_2\pi i}; q\right) + r_{im}(n|7)\right\}, \tag{365}$$

*and*

$$|r_{im}(n|7)| \le 54\theta_3(|z|^2 q^{2\beta_1}; q)\left\{\frac{\log n}{n^\rho} + \frac{q^{k_n}}{1-q} + |z|^{2j_n} q^{j_n^2+2(\beta_1-1)j_n} + |z|^{-2j_n} q^{j_n^2-2(\beta_1+1)j_n}\right\} \tag{366}$$

*for $n$ sufficiently large.*

In the following corollaries we assume that

$$z = e^{\pi u}, \quad u \in \mathbb{R}. \tag{367}$$

If we let

$$q = \exp(-n^{-a}\pi), \quad 0 < a < \frac{1}{2}, \quad n \in \mathbb{N}, \tag{368}$$

then we have the following results for Ismail-Masson polynomials:

**Corollary 5.2.** (1) *If$\tau > 0$, then,*

$$\begin{aligned} & h_n\left(\sinh \pi\left(u + (\tau+1/2)n^{1-a}\right) \mid \exp(-\pi n^{-a})\right) \\ & \quad = \exp(n\pi u + (\tau+1/2)\pi n^{2-a})\left\{1 + \mathcal{O}(e^{-4\pi n^a})\right\}, \end{aligned} \tag{369}$$

*and*

$$\begin{aligned} & h_n\left(\sinh \pi(u + (\tau+1/2)n^{1-a}) \| \exp(-\pi n^{-a})\right) \\ & \quad = \frac{1}{\sqrt{\pi}} \frac{\exp\left(-n^{-a}\pi(n^a u + \tau n)^2\right)}{\exp \frac{\pi}{12}\left(3n^{1-a} + n^{-a} - n^a\right)}\left\{1 + \mathcal{O}(e^{-4\pi n^a})\right\} \end{aligned} \tag{370}$$

*as $n \to \infty$.*

(2) *Assume that $-\frac{1}{2} < \tau < 0$ and for some fixed real number $\lambda$ there are infinitely many positive integers $n$ such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}, \tag{371}$$

*then,*

$$\begin{aligned} & h_n\left(\sinh \pi\left(u + (\tau+1/2)n^{1-a}\right) \mid \exp(-\pi n^{-a})\right) \\ & = \frac{(-1)^{\tau n+\lambda}\sqrt{2}\left\{\cos \pi(n^a u - \lambda) + \mathcal{O}(e^{-2\pi n^a})\right\}}{\exp\left\{-\pi n^{-a}(n^a u + (\tau+1/2)n)^2 - \frac{\pi n^{2-a}}{4} + \frac{\pi n^a}{12} + \frac{\pi n^{-a}}{24}\right\}}, \end{aligned} \tag{372}$$

*and*

$$\begin{aligned} & h_n\left(\sinh \pi(u + (\tau+1/2)n^{1-a}) \| \exp(-\pi n^{-a})\right) \\ & \quad = \sqrt{\frac{2}{\pi}} \frac{(-1)^{\tau n+\lambda}\left\{\cos \pi(n^a u - \lambda) + \mathcal{O}(e^{-2\pi n^a})\right\}}{\exp\left\{\frac{n^{1-a}\pi}{4} + \frac{\pi n^{-a}}{8}\right\}} \end{aligned} \tag{373}$$

*as $n \to \infty$.*

If we choose that

$$q = \exp\left(-\frac{\pi}{\gamma \log n}\right), \quad n \geq 2, \tag{374}$$

then we have the following:

**Corollary 5.3.** (1) *If $\tau > 0$, then we have*

$$\begin{aligned} & h_n\left(\sinh \pi\left(u + \frac{(\tau+1/2)n}{\gamma \log n}\right) \mid \exp\left(-\frac{\pi}{\gamma \log n}\right)\right) \\ & \quad = \exp \pi\left(nu + \frac{(\tau+1/2)n^2}{\gamma \log n}\right)\left\{1 + \mathcal{O}\left(\exp(-2\pi\tau n/(\gamma \log n)\right)\right\}, \end{aligned} \tag{375}$$

*and*

$$h_n\left(\sinh\pi\left(u+\frac{(\tau+1/2)n}{\gamma\log n}\right)\|\exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \tag{376}$$
$$==\frac{n^{\gamma\pi/12}}{\sqrt{\pi}}\frac{\exp\left(-\frac{\pi}{\gamma\log n}(u\gamma\log n+n\tau)^2\right)}{\exp\left(\frac{n\pi}{4\gamma\log n}+\frac{\pi}{12\gamma\log n}\right)}\left\{1+\mathcal{O}(n^{-4\pi\gamma})\right\}$$

*for $n$ sufficiently large.*

(2) *Assume that $-\frac{1}{2}<\tau<0$ and for some fixed real number $\lambda$ there are infinitely many $m$ and $n>0$ such that*

$$-n\tau=m+\lambda,\quad m\in\mathbb{N}. \tag{377}$$

*Then,*

$$h_n\left(\sinh\pi\left(u+\frac{(\tau+1/2)n}{\gamma\log n}\right)|\exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \tag{378}$$
$$=\frac{\sqrt{2}\exp\left(\frac{\pi}{\gamma\log n}(u\gamma\log n+(\tau+1/2)n)^2\right)}{(-1)^{\lambda+\tau n}n^{\pi\gamma/12}\exp(\frac{\pi}{24\gamma\log n}-\frac{n^2\pi}{4\gamma\log n})}$$
$$\times\left\{\cos\pi(u\gamma\log n-\lambda)+\mathcal{O}(n^{-2\pi\gamma})\right\},$$

*and*

$$h_n\left(\sinh\pi\left(u+\frac{(\tau+1/2)n}{\gamma\log n}\right)\|\exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \tag{379}$$
$$=\sqrt{\frac{2}{\pi}}\frac{\left\{\cos\pi(u\gamma\log n-\lambda)+\mathcal{O}(n^{-2\pi\gamma})\right\}}{(-1)^{\lambda+\tau n}\exp(\frac{\pi}{8\gamma\log n}+\frac{n\pi}{4\gamma\log n})}$$

*for $n$ sufficiently large.*

(3) *Assume that $-\frac{1}{2}<\tau<0$, if for some fixed real numbers $\beta$, $\rho\geq 1$ and $\lambda$ there exist infinitely many positive integers $n$ such that*

$$-n\tau=m+\beta+a_n,\quad |a_n|<\frac{1}{n^\rho},\quad m\in\mathbb{N}, \tag{380}$$

*and let*

$$\gamma=\frac{4\rho}{9\pi}, \tag{381}$$

*then,*

$$h_n\left(\sinh\pi\left(u+\frac{(\tau+1/2)n}{\gamma\log n}\right)|\exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \tag{382}$$
$$=\frac{\sqrt{2}\exp\left(\frac{\pi}{\gamma\log n}(u\gamma\log n+(\tau+1/2)n)^2\right)}{(-1)^{m}n^{\pi\gamma/12}\exp(\frac{\pi}{24\gamma\log n}-\frac{n^2\pi}{4\gamma\log n})}$$
$$\times\left\{\cos\pi(u\gamma\log n-\beta)+\mathcal{O}(n^{-8\rho/9}\log^2 n)\right\},$$

*and*

$$\begin{aligned} & h_n\left(\sinh\pi\left(u+\frac{(\tau+1/2)n}{\gamma\log n}\right)\|\exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \\ &= \sqrt{\frac{2}{\pi}}\frac{\left\{\cos\pi(u\gamma\log n-\beta)+\mathcal{O}(n^{-8\rho/9}\log^2 n)\right\}}{(-1)^m\exp(\frac{\pi}{8\gamma\log n}+\frac{n\pi}{4\gamma\log n})}\end{aligned}\tag{383}$$

*for $n$ sufficiently large.*

5.2. **Proof For Theorem 5.1.** In first three proofs we have used the following inequalities

$$0<\frac{(q;q)_n}{(q;q)_{n-k}}\le 1\tag{384}$$

for $0\le k\le n$ and

$$\left|\frac{(q;q)_n}{(q;q)_{n-k}}-1\right|\le\frac{6q^{1+n/2}}{1-q}\tag{385}$$

for $0\le k\le\left\lfloor\frac{n}{2}\right\rfloor-1$ and $n$ sufficiently large. This can be seen by applying Lemma 1.1

$$\frac{(q;q)_n}{(q;q)_{n-k}}-1=\{r_1(q;n-k)+1\}\{r_2(q;n)+1\}-1.\tag{386}$$

Assume that

$$-\tau n=m+c_n,\quad m\in\mathbb{N},\quad n\theta=m_1+d_n,\quad m_1\in\mathbb{Z},\tag{387}$$

then,

$$\begin{aligned}\frac{h_n(\sinh_n|q)}{z^nq^{-n^2s}} &= \sum_{k=0}^{n}\frac{q^{k^2}(q;q)_n e^{2nk\theta\pi i}}{(q;q)_k(q;q)_{n-k}}\left(-\frac{q^{2\tau n}}{z^2}\right)^k \\ &= \sum_{k=0}^{m}\frac{q^{k^2}(q;q)_n e^{2nk\theta\pi i}}{(q;q)_k(q;q)_{n-k}}\left(-\frac{q^{2\tau n}}{z^2}\right)^k+\sum_{k=m+1}^{n}\frac{q^{k^2}(q;q)_n e^{2nk\theta\pi i}}{(q;q)_k(q;q)_{n-k}}\left(-\frac{q^{2\tau n}}{z^2}\right)^k \\ &= s_1+s_2.\end{aligned}\tag{388}$$

We reverse the summation order in $s_1$ to obtain

$$\frac{s_1(q;q)_\infty(-z^2e^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}}=\sum_{k=0}^{m}q^{k^2}\left(-z^2q^{2c_n}e^{-2\pi i d_n}\right)^k e(k,n),\tag{389}$$

and

$$e(k,n)=\frac{(q;q)_\infty(q;q)_n}{(q;q)_{m-k}(q;q)_{n-m+k}}.\tag{390}$$

It is clear that

$$|e(k,n)|\le 1\tag{391}$$

for $0\le k\le m$. Expand

$$e(k,n)-1=\{r_2\,(q;n)+1\}\{r_1\,(q;m-k)+1\}\{r_1\,(q;n-m+k)+1\}-1\tag{392}$$

and estimate each term by Lemma 1.1 to get

$$|e(k,n)-1|\le\frac{14q^{2+k_n}}{1-q}\tag{393}$$

for $0\le k\le k_n-1$ and $n$ sufficiently large.

In sum $s_2$ we shift summation index from $k$ to $k+m$ to get

$$\frac{s_2(q;q)_\infty(-z^2e^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}} = \sum_{k=1}^{n-m} q^{k^2}\left(-z^{-2}q^{-2c_n}e^{2\pi i d_n}\right)^k f(k,n), \tag{394}$$

and

$$f(k,n) = \frac{(q;q)_\infty(q;q)_n}{(q;q)_{m+k}(q;q)_{n-m-k}}, \quad |f(k,n)| \le 1, \tag{395}$$

for $1 \le k \le n-m$. Apply Lemma 1.1 to each term of

$$f(k,n)-1 = \{r_2(q;n)+1\}\{r_1(q;m+k)+1\}\{r_1(q;n-m-k)+1\}-1, \tag{396}$$

to obtain

$$|f(k,n)-1| \le \frac{14q^{2+k_n}}{1-q} \tag{397}$$

for $1 \le k \le k_n - 1$ for $n$ sufficiently large.

5.2.1. *Proof for case 1.* If we write

$$\frac{h_n(\sinh\xi_n|q)}{z^nq^{-n^2s}} = 1 + r_{im}(n|1), \tag{398}$$

then

$$r_{im}(n|1) = \sum_{k=1}^n \frac{q^{k^2}(q;q)_n e^{2nk\theta\pi i}}{(q;q)_k(q;q)_{n-k}}\left(-\frac{q^{2\tau n}}{z^2}\right)^k. \tag{399}$$

Thus

$$|r_{im}(n|1)| \le \sum_{k=1}^\infty \frac{q^{k^2}}{(q;q)_k}\left(\frac{q^{2\tau n}}{|z|^2}\right)^k \le \sum_{k=1}^\infty \frac{1}{k!}\left(\frac{q^{2\tau n+1}}{|z|^2(1-q)}\right)^k \le \frac{q^{2\tau n+1}}{|z|^2(1-q)}\exp\left(\frac{q^{2\tau n+1}}{|z|^2(1-q)}\right). \tag{400}$$

5.2.2. *Proof for case 2.* We have

$$\begin{aligned} \frac{h_n(\sinh\xi_n|q)}{z^nq^{-n^2s}} &= \sum_{k=0}^n \frac{q^{k^2}}{(q;q)_k}\left(-\frac{e^{2\lambda\pi i}}{z^2}\right)^k \frac{(q;q)_n}{(q;q)_{n-k}} \\ &= \sum_{k=0}^\infty \frac{q^{k^2}}{(q;q)_k}\left(-\frac{e^{2\lambda\pi i}}{z^2}\right)^k - \sum_{k=\lfloor n/2\rfloor}^\infty \frac{q^{k^2}}{(q;q)_k}\left(-\frac{e^{2\lambda\pi i}}{z^2}\right)^k \\ &+ \sum_{k=0}^{\lfloor n/2\rfloor-1} \frac{q^{k^2}}{(q;q)_k}\left(-\frac{e^{2\lambda\pi i}}{z^2}\right)^k\left\{\frac{(q;q)_n}{(q;q)_{n-k}}-1\right\} + \sum_{k=\lfloor n/2\rfloor}^n \frac{q^{k^2}}{(q;q)_k}\left(-\frac{e^{2\lambda\pi i}}{z^2}\right)^k \frac{(q;q)_n}{(q;q)_{n-k}} \\ &= A_q\left(\frac{e^{2\lambda\pi i}}{z^2}\right) + s_1 + s_2 + s_3. \end{aligned} \tag{401}$$

Then,

$$|s_1+s_3| \le 2\sum_{k=\lfloor n/2\rfloor}^\infty \frac{q^{k^2}}{(q;q)_k}\left(\frac{1}{|z|^2}\right)^k \le \frac{2q^{\lfloor n/2\rfloor^2}A_q(-|z|^{-2})}{(q;q)_\infty|z|^{2\lfloor n/2\rfloor}}, \tag{402}$$

and

$$|s_2| \le \frac{6q^{n/2}}{1-q} A_q(-|z|^{-2}). \tag{403}$$

for $n$ sufficiently large.

Let

$$r_{im}(n|2) = s_1 + s_2 + s_3, \tag{404}$$

then,

$$\frac{h_n(\sinh \xi_n|q)}{z^n q^{-n^2 s}} = A_q\left(\frac{e^{2\lambda\pi i}}{z^2}\right) + r_{im}(n|2), \tag{405}$$

and

$$|r_{im}(n|2)| \le 6\frac{A_q(-|z|^{-2})}{(q;q)_\infty}\left\{q^{n/2} + \frac{q^{\lfloor n/2\rfloor^2}}{|z|^{2\lfloor n/2\rfloor}}\right\}. \tag{406}$$

5.2.3. *Proof for case 3.* In this case we have

(407)

$$\begin{aligned}
\frac{h_n(\sinh \xi_n|q)}{z^n q^{-n^2 s}} &= \sum_{k=0}^{n} \frac{q^{k^2}}{(q;q)_k}\left(-\frac{e^{2\pi i\beta}}{z^2}\right)^k \frac{(q;q)_n e^{2\pi i k b_n}}{(q;q)_{n-k}} \\
&= \sum_{k=0}^{\infty} \frac{q^{k^2}}{(q;q)_k}\left(-\frac{e^{2\pi i\beta}}{z^2}\right)^k - \sum_{k=j_n}^{\infty} \frac{q^{k^2}}{(q;q)_k}\left(-\frac{e^{2\pi i\beta}}{z^2}\right)^k \\
&+ \sum_{k=0}^{j_n-1} \frac{q^{k^2}}{(q;q)_k}\left(-\frac{e^{2\pi i\beta}}{z^2}\right)^k \left\{\frac{(q;q)_n}{(q;q)_{n-k}} - 1\right\} + \sum_{k=0}^{j_n-1} \frac{q^{k^2}}{(q;q)_k}\left(-\frac{e^{2\pi i\beta}}{z^2}\right)^k \frac{(q;q)_n}{(q;q)_{n-k}}\left\{e^{2\pi i k b_n} - 1\right\} \\
&+ \sum_{k=j_n}^{n} \frac{q^{k^2}}{(q;q)_k}\left(-\frac{e^{2\pi i\beta}}{z^2}\right)^k \frac{(q;q)_n}{(q;q)_{n-k}} e^{2\pi i k b_n} = A_q\left(\frac{e^{2\pi i\beta}}{z^2}\right) + s_1 + s_2 + s_3 + s_4.
\end{aligned}$$

Then,

$$|s_1 + s_4| \le 2\sum_{k=j_n}^{\infty} \frac{q^{k^2}|z|^{-2k}}{(q;q)_k} \le \frac{2}{j_n!}\left(\frac{q}{|z|^2(1-q)}\right)^{j_n} \exp\left(\frac{q}{|z|^2(1-q)}\right). \tag{408}$$

Given a positive number $\rho$, since that

$$1 < j_n < \frac{n}{2}, \quad j_n \le \frac{n^\rho}{2\pi} \tag{409}$$

for $n$ sufficiently large, clearly,

$$|s_2| \le \frac{6q^{n/2}}{1-q} \exp\left(\frac{q}{|z|^2(1-q)}\right), \tag{410}$$

and

$$|s_3| \le \frac{2\pi j_n}{n^\rho} e^{2\pi j_n/n^\rho} \sum_{k=0}^{\infty} \frac{q^{k^2}|z|^{-2k}}{(q;q)_k} \le \frac{24\log n}{n^\rho} \exp\left(\frac{q}{|z|^2(1-q)}\right). \tag{411}$$

Let

$$e_{im}(n|3) = s_1 + s_2 + s_3 + s_4, \tag{412}$$

then

$$\frac{h_n(\sinh\xi_n|q)}{z^n q^{-n^2 s}} = A_q\left(\frac{e^{2\pi i\beta}}{z^2}\right) + e_{im}(n|3), \tag{413}$$

and

$$|e_{im}(n|3)| \le 24\exp\left(\frac{q}{|z|^2(1-q)}\right)\left\{\frac{\log n}{n^\rho} + \frac{q^{n/2}}{1-q} + \frac{(|z|^2(q^{-1}-1))^{-j_n}}{j_n!}\right\} \tag{414}$$

for $n$ sufficiently large.

5.2.4. *Proof for case 4.* Observe that

(415)

$$\frac{s_1\left(-z^2e^{-2\pi i\lambda_1}\right)^m (q;q)_\infty}{q^{m(2\tau n+m)}} = \sum_{k=0}^{\infty} q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\lambda_1}\right)^k$$
$$- \sum_{k=k_n}^{\infty} q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\lambda_1}\right)^k + \sum_{k=0}^{k_n-1} q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\lambda_1}\right)^k (e(k,n)-1)$$
$$+ \sum_{k=k_n}^{m} q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\lambda_1}\right)^k e(k,n) = \sum_{k=0}^{\infty} q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\lambda_1}\right)^k + s_{11} + s_{12} + s_{13}.$$

Then,

$$\begin{aligned}|s_{11}+s_{13}| &\le 2\sum_{k=k_n}^{\infty} q^{k^2}\left(|z|^2 q^{2\lambda}\right)^k \le 2|z|^{2k_n} q^{k_n^2+2\lambda k_n}\sum_{k=0}^{\infty} q^{k^2}\left(|z|^2 q^{2\lambda}\right)^k \\ &\le 2\theta_3\left(|z|^2 q^{2\lambda} \mid q\right)|z|^{2k_n}q^{k_n^2+2\lambda k_n},\end{aligned} \tag{416}$$

and

$$|s_{12}| \le \frac{14q^{k_n}}{1-q}\sum_{k=0}^{\infty} q^{k^2}(|z|^2q^{2\lambda})^k \le 14\theta_3\left(|z|^2 q^{2\lambda};q\right)\frac{q^{k_n}}{1-q} \tag{417}$$

for $n$ sufficiently large.

Let

$$r_1(n) = s_{11} + s_{12} + s_{13}, \tag{418}$$

then,

$$\frac{s_1\left(-z^2e^{-2\pi i\lambda_1}\right)^m (q;q)_\infty}{q^{m(2\tau n+m)}} = \sum_{k=0}^{\infty} q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\lambda_1}\right)^k + r_1(n) \tag{419}$$

$$|r_1(n)| \le 14\theta_3\left(|z|^2 q^{2\lambda};q\right)\left\{|z|^{2k_n}q^{k_n^2+2\lambda k_n} + \frac{q^{k_n}}{1-q}\right\} \tag{420}$$

for $n$ sufficiently large.

Similarly,

$$\begin{aligned} &\frac{s_2\left(-z^2e^{-2\pi i\lambda_1}\right)^m (q;q)_\infty}{q^{m(2\tau n+m)}} = \sum_{k=1}^{\infty} q^{k^2}\left(-z^{-2}q^{-2\lambda}e^{2\pi i\lambda_1}\right)^k \\ &- \sum_{k=k_n}^{\infty} q^{k^2}\left(-z^{-2}q^{-2\lambda}e^{2\pi i\lambda_1}\right)^k + \sum_{k=1}^{k_n-1} q^{k^2}\left(-z^{-2}q^{-2\lambda}e^{2\pi i\lambda_1}\right)^k (f(k,n)-1) \\ &+ \sum_{k=k_n}^{n-m} q^{k^2}\left(-z^{-2}q^{-2\lambda}e^{2\pi i\lambda_1}\right)^k f(k,n) = \sum_{k=-\infty}^{-1} q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\lambda_1}\right)^k + s_{21} + s_{22} + s_{23}. \end{aligned} \tag{421}$$

Then,

$$|s_{21}+s_{23}| \le 2\sum_{k=k_n}^{\infty} q^{k^2}\left(|z|^{-2}q^{-2\lambda}\right)^k \le 2\theta_3(|z|^2q^{2\lambda};q)\frac{q^{k_n^2-2\lambda k_n}}{|z|^{2k_n}}, \tag{422}$$

and

$$|s_{22}| \le \frac{14q^{k_n}}{1-q}\sum_{k=0}^{\infty} q^{k^2-2\lambda k}|z|^{-2k} \le 14\theta_3(|z|^2q^{2\lambda};q)\frac{q^{k_n}}{1-q}. \tag{423}$$

for $n$ sufficiently large.

Let

$$r_2(n) = s_{21} + s_{22} + s_{23}, \tag{424}$$

then,

$$\frac{s_2\left(-z^2e^{-2\pi i\lambda_1}\right)^m (q;q)_\infty}{q^{m(2\tau n+m)}} = \sum_{k=-\infty}^{-1} q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\lambda_1}\right)^k + r_2(n), \tag{425}$$

and

$$|r_2(n)| \le 14\theta_3\left(|z|^2q^{2\lambda};q\right)\left\{\frac{q^{k_n}}{1-q} + \frac{q^{k_n^2-2\lambda k_n}}{|z|^{2k_n}}\right\} \tag{426}$$

for $n$ sufficiently large.

Thus,

$$\frac{\left(-z^2e^{-2\pi i\lambda_1}\right)^m h_n(\sinh\xi_n|q)(q;q)_\infty}{z^nq^{-n^2s+m(2\tau n+m)}} = \theta_4\left(z^2q^{2\lambda}e^{-2\pi i\lambda_1};q\right) + r_{im}(n|4) \tag{427}$$

with

$$|r_{im}(n|4)| \le 28\theta_3\left(|z|^2q^{2\lambda};q\right)\left\{\frac{q^{k_n}}{1-q} + |z|^{2k_n}q^{k_n^2+2\lambda k_n} + |z|^{-2k_n}q^{k_n^2-2\lambda k_n}\right\} \tag{428}$$

for $n$ sufficiently large.

5.2.5. *Proof for case 5.* In this case we have

(429)

$$\begin{aligned}&\frac{s_1(q;q)_\infty(-z^2e^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}}=\sum_{k=0}^{m}q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\beta}\right)^k e^{-2k\pi i b_n}e(k,n)\\ &=\sum_{k=0}^{\infty}q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\beta}\right)^k-\sum_{k=j_n}^{\infty}q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\beta}\right)^k\\ &+\sum_{k=0}^{j_n-1}q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\beta}\right)^k\left\{e^{-2k\pi i b_n}-1\right\}+\sum_{k=0}^{j_n-1}q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\beta}\right)^k e^{-2k\pi i b_n}\left\{e(k,n)-1\right\}\\ &+\sum_{k=j_n}^{m}q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\beta}\right)^k e^{-2k\pi i b_n}e(k,n)=\sum_{k=0}^{\infty}q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\beta}\right)^k+s_{11}+s_{12}+s_{13}+s_{14}.\end{aligned}$$

Then,

(430) $$|s_{11}+s_{14}|\le 2\sum_{k=j_n}^{\infty}q^{k^2+2\lambda k}|z|^{2k}\le 2\theta_3(|z|^2q^{2\lambda}\mid q)|z|^{2j_n}q^{j_n^2+2\lambda j_n}.$$

For sufficiently large $n$,

(431) $$|s_{12}|\le\frac{24j_n}{n^\rho}\sum_{k=0}^{\infty}q^{k^2+2\lambda k}|z|^{2k}\le 24\theta_3(|z|^2q^{2\lambda}\mid q)\frac{\log n}{n^\rho},$$

and

(432) $$|s_{13}|\le\frac{14q^{k_n}}{1-q}\sum_{k=0}^{\infty}q^{k^2+2\lambda k}|z|^{2k}\le 14\theta_3(|z|^2q^{2\lambda}\mid q)\frac{q^{k_n}}{1-q}.$$

Let

(433) $$r_1(n)=s_{11}+s_{12}+s_{13}+s_{14},$$

then,

(434) $$\frac{s_1(q;q)_\infty(-z^2e^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}}=\sum_{k=0}^{\infty}q^{k^2}\left(-z^2q^{2\lambda}e^{-2\pi i\beta}\right)^k+r_1(n),$$

and

(435) $$|r_1(n)|\le 24\theta_3(|z|^2q^{2\lambda}\mid q)\left\{|z|^{2j_n}q^{j_n^2+2\lambda j_n}+\frac{q^{k_n}}{1-q}+\frac{\log n}{n^\rho}\right\}$$

for $n$ sufficiently large.

Similarly,

(436)

$$\begin{aligned}&\frac{s_2(q;q)_\infty(-z^2e^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}} = \sum_{k=1}^{n-m} q^{k^2}\left(-z^{-2}q^{-2\lambda}e^{2\beta\pi i}\right)^k e^{2k\pi i b_n} f(k,n)\\
&= \sum_{k=1}^{\infty} q^{k^2}\left(-z^{-2}q^{-2\lambda}e^{2\beta\pi i}\right)^k - \sum_{k=j_n}^{\infty} q^{k^2}\left(-z^{-2}q^{-2\lambda}e^{2\beta\pi i}\right)^k\\
&+ \sum_{k=1}^{j_n-1} q^{k^2}\left(-z^{-2}q^{-2\lambda}e^{2\beta\pi i}\right)^k\left\{e^{2k\pi i b_n}-1\right\} + \sum_{k=1}^{j_n-1} q^{k^2}\left(-z^{-2}q^{-2\lambda}e^{2\beta\pi i}\right)^k e^{2k\pi i b_n}\left\{f(k,n)-1\right\}\\
&+ \sum_{k=j_n}^{n-m} q^{k^2}\left(-z^{-2}q^{-2\lambda}e^{2\beta\pi i}\right)^k e^{2k\pi i b_n} f(k,n) = \sum_{k=-\infty}^{-1} q^{k^2}\left(-z^{2}q^{2\lambda}e^{-2\beta\pi i}\right)^k + s_{21}+s_{22}+s_{23}+s_{24},\end{aligned}$$

then

$$|s_{21}+s_{24}| \le 2\sum_{k=j_n}^{\infty} q^{k^2-2\lambda k}|z|^{-2k} \le 2\theta_3(|z|^2q^{2\lambda};q)\frac{q^{j_n^2-2\lambda j_n}}{|z|^{2j_n}}. \tag{437}$$

For sufficiently large $n$

$$|s_{22}| \le \frac{24j_n}{n^\rho}\sum_{k=0}^{\infty} q^{k^2-2\lambda k}|z|^{-2k} \le 24\theta_3(|z|^2q^{2\lambda};q)\frac{\log n}{n^\rho}, \tag{438}$$

and

$$|s_{23}| \le 14\frac{q^{k_n}}{1-q}\sum_{k=0}^{\infty} q^{k^2-2\lambda k}|z|^{-2k} \le 14\theta_3(|z|^2q^{2\lambda};q)\frac{q^{k_n}}{1-q}. \tag{439}$$

for $n$ sufficiently large.

Let

$$r_2(n) = s_{21}+s_{22}+s_{23}+s_{24}, \tag{440}$$

then,

$$\frac{s_2(q;q)_\infty(-z^2e^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}} = \sum_{k=-\infty}^{-1} q^{k^2}\left(-z^2q^{2\lambda}e^{-2\beta\pi i}\right)^k + r_2(n), \tag{441}$$

and

$$|r_2(n)| \le 24\theta_3(|z|^2q^{2\lambda};q)\left\{\frac{q^{j_n^2-2\lambda j_n}}{|z|^{2j_n}} + \frac{q^{k_n}}{1-q} + \frac{\log n}{n^\rho}\right\} \tag{442}$$

for $n$ sufficiently large.

Thus,

$$\frac{(-z^2e^{-2n\theta\pi i})^m h_n(\sinh\xi_n|q)(q;q)_\infty}{z^n q^{-n^2 s+m(2\tau n+m)}} = \theta_4\left(z^2q^{2\lambda}e^{-2\beta\pi i};q\right) + r_{im}(n|5), \tag{443}$$

and

$$|r_{im}(n|5)| \le 48\theta_3(|z|^2q^{2\lambda};q)\left\{\frac{q^{j_n^2-2\lambda j_n}}{|z|^{2j_n}} + |z|^{2j_n}q^{j_n^2+2\lambda j_n} + \frac{q^{k_n}}{1-q} + \frac{\log n}{n^\rho}\right\} \tag{444}$$

for sufficiently large $n$.

5.2.6. *Proof for case 6.* Observe that

(445)
$$\begin{aligned}
&\frac{s_1(q;q)_\infty(-z^2e^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}} = \sum_{k=0}^{m} q^{k^2}\left(-z^2q^{2\beta}e^{-2\pi i\lambda}\right)^k q^{2ka_n}e(k,n)\\
&= \sum_{k=0}^{\infty} q^{k^2}\left(-z^2q^{2\beta}e^{-2\pi i\lambda}\right)^k - \sum_{k=j_n}^{\infty} q^{k^2}\left(-z^2q^{2\beta}e^{-2\pi i\lambda}\right)^k\\
&+ \sum_{k=0}^{j_n-1} q^{k^2}\left(-z^2q^{2\beta}e^{-2\pi i\lambda}\right)^k\left\{q^{2ka_n}-1\right\} + \sum_{k=0}^{j_n-1} q^{k^2}\left(-z^2q^{2\beta}e^{-2\pi i\lambda}\right)^k q^{2ka_n}\left\{e(k,n)-1\right\}\\
&+ \sum_{k=j_n}^{m} q^{k^2}\left(-z^2q^{2\beta}e^{-2\pi i\lambda}\right)^k q^{2ka_n}e(k,n) = \sum_{k=0}^{\infty} q^{k^2}\left(-z^2q^{2\beta}e^{-2\pi i\lambda}\right)^k + s_{11}+s_{12}+s_{13}+s_{14}.
\end{aligned}$$

Since

$$|q^{2ka_n}| \le q^{-2k} \tag{446}$$

for $0 \le k \le n$, then,

$$|s_{11}+s_{14}| \le 2\sum_{k=j_n}^{\infty} q^{k^2}(|z|^2q^{2\beta-2})^k \le 2\theta_3(|z|^2q^{2\beta};q)|z|^{2j_n}q^{j_n^2+2(\beta-1)j_n}. \tag{447}$$

For $n$ sufficiently large we have

$$|q^{2ka_n}-1| \le \frac{6\log n}{n^\rho}, \quad |q^{2ka_n}| \le 3 \tag{448}$$

for $0 \le k \le j_n - 1$, then,

$$|s_{12}| \le 6\theta_3(|z|^2q^{2\beta};q)\frac{\log n}{n^\rho}, \tag{449}$$

and

$$|s_{13}| \le 3\theta_3(|z|^2q^{2\beta};q)\frac{q^{k_n}}{1-q}. \tag{450}$$

Thus,

$$\frac{s_1(q;q)_\infty(-z^2e^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}} = \sum_{k=0}^{\infty} q^{k^2}\left(-z^2q^{2\beta}e^{-2\pi i\lambda}\right)^k + r_1(n), \tag{451}$$

and

$$|r_1(n)| \le 6\theta_3(|z|^2q^{2\beta};q)\left\{\frac{\log n}{n^\rho} + \frac{q^{k_n}}{1-q} + |z|^{2j_n}q^{j_n^2+2(\beta-1)j_n}\right\} \tag{452}$$

for $n$ sufficiently large.

Similarly,

$$\begin{aligned}\frac{s_2(q;q)_\infty(-z^2e^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}} &= \sum_{k=1}^{n-m} q^{k^2}\left(-z^{-2}q^{-2\beta}e^{2\lambda\pi i}\right)^k q^{-2ka_n}f(k,n)\\ &= \sum_{k=-\infty}^{-1} q^{k^2}\left(-z^2q^{2\beta}e^{-2\pi i\lambda}\right)^k - \sum_{k=j_n}^{\infty} q^{k^2}\left(-z^{-2}q^{-2\beta}e^{2\lambda\pi i}\right)^k\\ &+ \sum_{k=1}^{j_n-1} q^{k^2}\left(-z^{-2}q^{-2\beta}e^{2\lambda\pi i}\right)^k\left\{q^{-2ka_n}-1\right\} + \sum_{k=1}^{j_n-1} q^{k^2}\left(-z^{-2}q^{-2\beta}e^{2\lambda\pi i}\right)^k q^{-2ka_n}\left\{f(k,n)-1\right\}\\ &+ \sum_{k=j_n}^{n-m} q^{k^2}\left(-z^{-2}q^{-2\beta}e^{2\lambda\pi i}\right)^k q^{-2ka_n}f(k,n) = \sum_{k=-\infty}^{-1} q^{k^2}\left(-z^2q^{2\beta}e^{-2\pi i\lambda}\right)^k + s_{21}+s_{22}+s_{23}+s_{24}.\end{aligned}\tag{453}$$

Then,

$$|s_{21}+s_{24}| \le 2\sum_{k=j_n}^{\infty} q^{k^2}(|z|^{-2}q^{-2\beta-2})^k \le 2\theta_3(|z|^2q^{2\beta};q)|z|^{-2j_n}q^{j_n^2-2(\beta+1)j_n}.\tag{454}$$

For $n$ sufficiently large,

$$|s_{22}| \le 6\theta_3(|z|^2q^{2\beta};q)\frac{\log n}{n^\rho},\tag{455}$$

and

$$|s_{23}| \le 3\theta_3(|z|^2q^{2\beta};q)\frac{q^{k_n}}{1-q}.\tag{456}$$

Therefore,

$$\frac{s_2(q;q)_\infty(-z^2e^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}} = \sum_{k=-\infty}^{-1} q^{k^2}\left(-z^2q^{2\beta}e^{-2\pi i\lambda}\right)^k + r_2(n),\tag{457}$$

and

$$|r_2(n)| \le 6\theta_3(|z|^2q^{2\beta};q)\left\{\frac{\log n}{n^\rho}+\frac{q^{k_n}}{1-q}+|z|^{-2j_n}q^{j_n^2-2(\beta+1)j_n}\right\}\tag{458}$$

for $n$ sufficiently large.

Hence,

$$\frac{(-z^2e^{-2n\theta\pi i})^m h_n(\sinh\xi_n|q)(q;q)_\infty}{z^nq^{-n^2s+m(2\tau n+m)}} = \theta_4(z^2q^{2\beta}e^{-2\pi i\lambda};q)+r_{im}(n|6),\tag{459}$$

and

$$|r_{im}(n|6)| \le 12\theta_3(|z|^2q^{2\beta};q)\left\{\frac{\log n}{n^\rho}+\frac{q^{k_n}}{1-q}+|z|^{2j_n}q^{j_n^2+2(\beta-1)j_n}+|z|^{-2j_n}q^{j_n^2-2(\beta+1)j_n}\right\}\tag{460}$$

for $n$ sufficiently large.

5.2.7. *Proof for case 7.* In this case we have

(461)

$$\frac{s_1(q;q)_\infty(-z^2e^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}}=\sum_{k=0}^{m}q^{k^2}\left(-z^2q^{2\beta_1}e^{-2\pi i\beta_2}\right)^k e^{-2k\pi ib_n}q^{2ka_n}e(k,n)$$

$$=\sum_{k=0}^{\infty}q^{k^2}\left(-z^2q^{2\beta_1}e^{-2\pi i\beta_2}\right)^k-\sum_{k=j_n}^{\infty}q^{k^2}\left(-z^2q^{2\beta_1}e^{-2\pi i\beta_2}\right)^k$$

$$+\sum_{k=0}^{j_n-1}q^{k^2}\left(-z^2q^{2\beta_1}e^{-2\pi i\beta_2}\right)^k\left\{q^{2ka_n}-1\right\}+\sum_{k=0}^{j_n-1}q^{k^2}\left(-z^2q^{2\beta_1}e^{-2\pi i\beta_2}\right)^k\left\{e^{-2k\pi ib_n}-1\right\}q^{2ka_n}$$

$$+\sum_{k=0}^{j_n-1}q^{k^2}\left(-z^2q^{2\beta_1}e^{-2\pi i\beta_2}\right)^k e^{-2k\pi ib_n}q^{2ka_n}\left\{e(k,n)-1\right\}+\sum_{k=j_n}^{m}q^{k^2}\left(-z^2q^{2\beta_1}e^{-2\pi i\beta_2}\right)^k e^{-2k\pi ib_n}q^{2ka_n}e(k,n)$$

$$=\sum_{k=0}^{\infty}q^{k^2}\left(-z^2q^{2\beta_1}e^{-2\pi i\beta_2}\right)^k+s_{11}+s_{12}+s_{13}+s_{14}+s_{15}.$$

Then,

$$\frac{s_1(q;q)_\infty(-z^2e^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}}=\sum_{k=0}^{\infty}q^{k^2}\left(-z^2q^{2\beta_1}e^{-2\pi i\beta_2}\right)^k+r_1(n), \tag{462}$$

and

$$|r_1(n)|\le 27\theta_3(|z|^2q^{2\beta_1};q)\left\{\frac{\log n}{n^\rho}+\frac{q^{k_n}}{1-q}+|z|^{2j_n}q^{j_n^2+2(\beta_1-1)j_n}\right\} \tag{463}$$

for $n$ sufficiently large.

Similarly,

(464)

$$\frac{s_2(q;q)_\infty(-z^2e^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}}=\sum_{k=1}^{n-m}q^{k^2}\left(-z^{-2}q^{-2\beta_1}e^{2\beta_2\pi i}\right)^k e^{2k\pi ib_n}q^{-2ka_n}f(k,n)$$

$$=\sum_{k=-\infty}^{-1}q^{k^2}\left(-z^2q^{2\beta_1}e^{-2\beta_2\pi i}\right)^k-\sum_{k=j_n}^{\infty}q^{k^2}\left(-z^{-2}q^{-2\beta_1}e^{2\beta_2\pi i}\right)^k$$

$$+\sum_{k=1}^{j_n-1}q^{k^2}\left(-z^{-2}q^{-2\beta_1}e^{2\beta_2\pi i}\right)^k\left\{e^{2k\pi ib_n}-1\right\}+\sum_{k=1}^{j_n-1}q^{k^2}\left(-z^{-2}q^{-2\beta_1}e^{2\beta_2\pi i}\right)^k e^{2k\pi ib_n}\left\{q^{-2ka_n}-1\right\}$$

$$+\sum_{k=1}^{j_n-1}q^{k^2}\left(-z^{-2}q^{-2\beta_1}e^{2\beta_2\pi i}\right)^k e^{2k\pi ib_n}q^{-2ka_n}\left\{f(k,n)-1\right\}+\sum_{k=j_n}^{n-m}q^{k^2}\left(-z^{-2}q^{-2\beta_1}e^{2\beta_2\pi i}\right)^k e^{2k\pi ib_n}q^{-2ka_n}f(k,n)$$

$$=\sum_{k=-\infty}^{-1}q^{k^2}\left(-z^2q^{2\beta_1}e^{-2\beta_2\pi i}\right)^k+s_{21}+s_{22}+s_{23}+s_{24}+s_{25}.$$

Thus,

$$\frac{s_2(q;q)_\infty(-z^2e^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}}=\sum_{k=-\infty}^{-1}q^{k^2}\left(-z^2q^{2\beta_1}e^{-2\beta_2\pi i}\right)^k+r_2(n), \tag{465}$$

and

$$|r_2(n)| \le 27\theta_3(|z|^2q^{2\beta_1};q)\left\{\frac{\log n}{n^\rho}+\frac{q^{k_n}}{1-q}+|z|^{-2j_n}q^{j_n^2-2(\beta_1+1)j_n}\right\} \tag{466}$$

for $n$ sufficiently large.

Therefore,

$$\frac{(-z^2e^{-2n\theta\pi i})^m h_n(\sinh\xi_n|q)(q;q)_\infty}{z^nq^{-n^2s+m(2\tau n+m)}} = \theta_4(z^2q^{2\beta_1}e^{-2\pi i\beta_2};q)+r_{im}(n|7), \tag{467}$$

and

(468)

$$|r_{im}(n|7)| \le 54\theta_3(|z|^2q^{2\beta_1};q)\left\{\frac{\log n}{n^\rho}+\frac{q^{k_n}}{1-q}+|z|^{2j_n}q^{j_n^2+2(\beta_1-1)j_n}+|z|^{-2j_n}q^{j_n^2-2(\beta_1+1)j_n}\right\}$$

For $n$ sufficiently large.

5.3. **Proof for Corollary 5.2.** In this case we have,

$$x_n(u)=\sinh\xi_n=\sinh\pi\left(u+(\tau+1/2)n^{1-a}\right), \tag{469}$$

and

$$w_{im}(x_n(u))=\frac{\sqrt{2n^a}}{\pi}\exp\left\{-\frac{n^{-a}\pi}{8}-2n^a\pi\left(u+(\tau+1/2)n^{1-a}\right)^2\right\}. \tag{470}$$

From Lemma 1.1 and Lemma 1.2 to get

$$\frac{1}{(q;q)_n}=\frac{r_1(q;n)+1}{(q;q)_\infty}=\frac{\exp(\pi n^a/6-\pi n^{-a}/24)}{\sqrt{2n^a}}\left\{1+\mathcal{O}(e^{-4\pi n^a})\right\} \tag{471}$$

as $n\to\infty$, thus,

$$\begin{aligned} q^{n(n+1)/4}\sqrt{\frac{w_{im}(x_n(u))}{(q;q)_n}} &= \frac{\exp\left\{-n^{-a}\pi(n^au+(\tau+1/2)n)^2+\frac{\pi n^a}{12}-\frac{\pi n^{-a}}{12}-\frac{(n+1)n\pi}{4}\right\}}{\sqrt{\pi}} \\ &\times \left\{1+\mathcal{O}(e^{-4\pi n^a})\right\} \end{aligned} \tag{472}$$

as $n\to\infty$.

5.3.1. *Proof of case 1.* Observe that

$$z^nq^{-n^2(\tau+1/2)}=\exp(n\pi u+(\tau+1/2)\pi n^{2-a}), \tag{473}$$

then formulas (344) and (345) imply

$$\begin{aligned} &h_n\left(\sinh\pi\left(u+(\tau+1/2)n^{1-a}\right)|\exp(-\pi n^{-a})\right) \\ &=\exp(n\pi u+(\tau+1/2)\pi n^{2-a})\left\{1+\mathcal{O}(e^{-4\pi n^a})\right\}, \end{aligned} \tag{474}$$

and

$$\begin{aligned} &h_n\left(\sinh\pi(u+(\tau+1/2)n^{1-a})\|\exp(-\pi n^{-a})\right) \\ &=\frac{1}{\sqrt{\pi}}\frac{\exp\left(-n^{-a}\pi(n^au+\tau n)^2\right)}{\exp\frac{\pi}{12}\left(3n^{1-a}+n^{-a}-n^a\right)}\left\{1+\mathcal{O}(e^{-4\pi n^a})\right\} \end{aligned} \tag{475}$$

as $n\to\infty$.

5.3.2. *Proof for case 2.* From (19)– (32) to get

$$\begin{aligned}\theta_3(|z|^2q^{2\lambda};q) &= \theta_3(ui-\lambda n^{-a}i \mid n^{-a}i)\\ &= \sqrt{n^a}\exp \pi n^{-a}(n^a u-\lambda)^2\left\{1+\mathcal{O}(e^{-\pi n^a})\right\},\end{aligned}\tag{476}$$

and

$$\begin{aligned}\theta_4(z^2q^{2\lambda};q) &= \theta_4(ui-\lambda n^{-a}i \mid n^{-a}i)\\ &= \sqrt{n^a}\exp \pi n^{-a}(n^a u-\lambda)^2\theta_2\left(n^a u-\lambda|n^a i\right)\\ &= 2\sqrt{n^a}\exp\left\{\pi n^{-a}(n^a u-\lambda)^2-\frac{\pi n^a}{4}\right\}\\ &\times \cos\pi(n^a u-\lambda)\left\{1+\mathcal{O}(e^{-2\pi n^a})\right\}\end{aligned}\tag{477}$$

as $n\to\infty$.

From formulas (353)–(356), we have

$$\begin{aligned}&h_n\left(\sinh\pi\left(u+(\tau+1/2)n^{1-a}\right)|\exp(-\pi n^{-a})\right)\\ &=\frac{(-1)^{\tau n+\lambda}\sqrt{2}\left\{\cos\pi(n^a u-\lambda)+\mathcal{O}(e^{-2\pi n^a})\right\}}{\exp\left\{-\pi n^{-a}(n^a u+(\tau+1/2)n)^2-\frac{\pi n^{2-a}}{4}+\frac{\pi n^a}{12}+\frac{\pi n^{-a}}{24}\right\}}\end{aligned}\tag{478}$$

and

$$\begin{aligned}&h_n\left(\sinh\pi(u+(\tau+1/2)n^{1-a})\|\exp(-\pi n^{-a})\right)\\ &=\sqrt{\frac{2}{\pi}}\frac{(-1)^{\tau n+\lambda}\left\{\cos\pi(n^a u-\lambda)+\mathcal{O}(e^{-2\pi n^a})\right\}}{\exp\left\{\frac{n^{1-a}\pi}{4}+\frac{\pi n^{-a}}{8}\right\}}\end{aligned}\tag{479}$$

as $n\to\infty$.

5.4. **Proof for Corollary 5.3.** In this case we have

$$\sinh\xi_n=\sinh\pi\left(u+\frac{(\tau+1/2)n}{\gamma\log n}\right),\tag{480}$$

$$w_{im}(\sinh\xi_n)=\frac{\sqrt{2\gamma\log n}}{\pi}\exp\left(-\frac{\pi}{8\gamma\log n}-\frac{2\pi}{\gamma\log n}\left(\gamma u\log n+(\tau+1/2)n\right)^2\right),\tag{481}$$

and

$$\frac{1}{(q;q)_n}=\frac{\exp\left(\frac{\gamma\pi\log n}{6}-\frac{\pi}{24\gamma\log n}\right)}{\sqrt{2\gamma\log n}}\left\{1+\mathcal{O}(n^{-4\pi\gamma})\right\},\tag{482}$$

$$\begin{aligned}&q^{n(n+1)/4}\sqrt{\frac{w_{im}(\sinh\xi_n)}{(q;q)_n}}\\ &=\frac{n^{\gamma\pi/12}\exp\left(-\frac{\pi}{\gamma\log n}\left(\gamma u\log n+(\tau+1/2)n\right)^2\right)}{\sqrt{\pi}\exp\left(\frac{\pi}{12\gamma\log n}+\frac{n(n+1)\pi}{4\gamma\log n}\right)}\left\{1+\mathcal{O}(n^{-4\pi\gamma})\right\}\end{aligned}\tag{483}$$

as $n\to\infty$.

5.4.1. *Proof for case 1.* From formulas (344) and (345) to get

$$\begin{aligned} h_n\left(\sinh\pi\left(u+\frac{(\tau+1/2)n}{\gamma\log n}\right) \mid \exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \\ = \exp\pi\left(nu+\frac{(\tau+1/2)n^2}{\gamma\log n}\right)\left\{1+\mathcal{O}\left(\exp(-2\pi\tau n/(\gamma\log n)\right)\right\}, \end{aligned} \tag{484}$$

and

$$\begin{aligned} h_n\left(\sinh\pi\left(u+\frac{(\tau+1/2)n}{\gamma\log n}\right) \| \exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \\ == \frac{n^{\gamma\pi/12}}{\sqrt{\pi}}\frac{\exp\left(-\frac{\pi(u\gamma\log n+n\tau)^2}{\gamma\log n}\right)}{\exp\left(\frac{n\pi}{4\gamma\log n}+\frac{\pi}{12\gamma\log n}\right)}\left\{1+\mathcal{O}(n^{-4\pi\gamma})\right\} \end{aligned} \tag{485}$$

as $n\to\infty$.

5.4.2. *Proof for case 2.* In this case we have

$$\begin{aligned} \theta_4(z^2q^{2\lambda};q) &= \theta_4\left(ui-\frac{\lambda i}{\gamma\log n} \mid \frac{i}{\gamma\log n}\right) \\ &= \frac{\sqrt{\gamma\log n}}{n^{\pi\gamma/4}}\exp\left(\frac{\pi(u\gamma\log n-\lambda)^2}{\gamma\log n}\right) \\ &\quad \times\cos\pi(u\gamma\log n-\lambda)\left\{1+\mathcal{O}(n^{-2\pi\gamma})\right\}, \end{aligned} \tag{486}$$

and

$$\begin{aligned} \theta_3(|z|^2q^{2\lambda};q) &= \theta_3\left(ui-\frac{\lambda i}{\gamma\log n} \mid \frac{i}{\gamma\log n}\right) \\ &= \sqrt{\gamma\log n}\exp\left(\frac{\pi(u\gamma\log n-\lambda)^2}{\gamma\log n}\right)\left\{1+\mathcal{O}(n^{-\pi\gamma})\right\}. \end{aligned} \tag{487}$$

Then,

$$\begin{aligned} h_n\left(\sinh\pi\left(u+\frac{(\tau+1/2)n}{\gamma\log n}\right) \mid \exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \\ = \frac{\sqrt{2}\exp\left(\frac{\pi}{\gamma\log n}(u\gamma\log n+(\tau+1/2)n)^2\right)}{(-1)^{\lambda+\tau n}n^{\pi\gamma/12}\exp(\frac{\pi}{24\gamma\log n}-\frac{n^2\pi}{4\gamma\log n})} \\ \times\left\{\cos\pi(u\gamma\log n-\lambda)+\mathcal{O}(n^{-2\pi\gamma})\right\}, \end{aligned} \tag{488}$$

and

$$\begin{aligned} h_n\left(\sinh\pi\left(u+\frac{(\tau+1/2)n}{\gamma\log n}\right) \| \exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \\ = \sqrt{\frac{2}{\pi}}\frac{\left\{\cos\pi(u\gamma\log n-\lambda)+\mathcal{O}(n^{-2\pi\gamma})\right\}}{(-1)^{\lambda+\tau n}\exp(\frac{\pi}{8\gamma\log n}+\frac{n\pi}{4\gamma\log n})} \end{aligned} \tag{489}$$

for $n$ sufficiently large.

5.4.3. *Proof for case 3.* Put $\theta, m_1, \lambda = 0$ in (359)-(361) to get

$$\begin{aligned} &h_n\left(\sinh\pi\left(u+\frac{(\tau+1/2)n}{\gamma\log n}\right) \,|\, \exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \\ &= \frac{\sqrt{2}\exp\left(\frac{\pi}{\gamma\log n}(u\gamma\log n+(\tau+1/2)n)^2\right)}{(-1)^m n^{\pi\gamma/12}\exp(\frac{\pi}{24\gamma\log n}-\frac{n^2\pi}{4\gamma\log n})} \\ &\times\left\{\cos\pi(u\gamma\log n-\beta)+\mathcal{O}(n^{-8\rho/9}\log^2 n)\right\}, \end{aligned} \tag{490}$$

and

$$\begin{aligned} &h_n\left(\sinh\pi\left(u+\frac{(\tau+1/2)n}{\gamma\log n}\right) \| \exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \\ &= \sqrt{\frac{2}{\pi}}\frac{\left\{\cos\pi(u\gamma\log n-\beta)+\mathcal{O}(n^{-8\rho/9}\log^2 n)\right\}}{(-1)^m\exp(\frac{\pi}{8\gamma\log n}+\frac{n\pi}{4\gamma\log n})} \end{aligned} \tag{491}$$

for $n$ sufficiently large, where $\gamma = \frac{8\rho}{9\pi}$.

## 6. Stieltjes-Wigert Orthogonal Polynomials

Stieltjes-Wigert orthogonal polynomials $\{S_n(x;q)\}_{n=0}^{\infty}$ are defined as [13]

$$S_n(x;q) = \sum_{k=0}^{n}\frac{q^{k^2}(-x)^k}{(q;q)_k(q;q)_{n-k}}. \tag{492}$$

Stieltjes-Wigert orthogonal polynomials come from an indeterminant moment problem. They satisfy the orthogonality relation

$$\int_0^{\infty} S_m(x;q)S_n(x;q)w_{sw}(x)dx = \frac{q^{-n}}{(q;q)_n}\delta_{m,n}, \tag{493}$$

where

$$w_{sw}(x) := \sqrt{\frac{-1}{2\pi\log q}}\exp\left(\frac{1}{2\log q}\left[\log\left(\frac{x}{\sqrt{q}}\right)\right]^2\right). \tag{494}$$

Clearly, the associated orthonormal Stieltjes-Wigert functions are given by

$$s_n(x\|q) := \sqrt{q^n(q;q)_n w_{sw}(x)}S_n(x;q). \tag{495}$$

For any nonzero complex number $z$, let

$$s = 2\tau+2+i\frac{2\theta\pi}{\log q}, \quad x_n(z,s) = zq^{-ns}, \quad \tau,\theta\in\mathbb{R}. \tag{496}$$

Reverse the summation order of (492) to get

$$\frac{S_n(x_n(z,s);q)}{(-z)^n q^{n^2(1-s)}} = \sum_{k=0}^{n}\frac{q^{k^2}e^{2nk\theta\pi i}}{(q;q)_k(q;q)_{n-k}}\left(-\frac{q^{2\tau n}}{z}\right)^k. \tag{497}$$

Clearly,

$$|S_n(x_n(z,s);q)| \le \frac{|z|^n q^{-n^2(2\tau+1)}}{(q;q)_\infty}\sum_{k=0}^{\infty}\frac{q^{k^2}(q^{2\tau n}|z|^{-1})^k}{(q;q)_k} \le \frac{|z|^n A_q(-q^{2\tau n}|z|^{-1})}{(q;q)_\infty q^{n^2(2\tau+1)}}, \tag{498}$$

or

$$|S_n(x_n(z,s);q)| \le \frac{(-\sqrt{q};q)_\infty}{(q;q)_\infty} \frac{|z|^{(2\tau+1)n} \exp\left(-\frac{\log^2|z|}{2\log q}\right)}{q^{(2\tau^2+2\tau+1)n^2}}. \tag{499}$$

## 6.1. **Asymptotic Formulas For Stieltjes-Wigert Polynomials.**

**Theorem 6.1.** *Given any nonzero complex number $z$, let $s$ and $x_n(z,s)$ be defined as in* (493) *and* (494) *and let*

$$j_n := \left\lfloor \frac{q^4 \log n}{\log q^{-1}} \right\rfloor, \quad k_n = \min\left\{ \left\lfloor \frac{(1+\tau)n}{2} \right\rfloor, \left\lfloor \frac{(-\tau)n}{2} \right\rfloor \right\}, \tag{500}$$

*we have the following results for Stieltjes-Wigert polynomials:*

1. *When $\tau > 0$, we have*

$$\frac{S_n(x_n(z,s);q)(q;q)_\infty}{(-z)^n q^{n^2(1-s)}} = 1 + r_{sw}(n|1), \tag{501}$$

*and*

$$|r_{sw}(n|1)| \le \frac{q^{2\tau n+1} \exp(\big(q^{2\tau n+1}/((1-q)|z|)\big)}{(1-q)|z|}. \tag{502}$$

2. *Assume that $\tau = 0$, if for a fixed real number $\lambda$ there are infinitely many positive integers $n$ such that*

$$n\theta = m + \lambda, \quad m \in \mathbb{Z}, \tag{503}$$

*then,*

$$\frac{S_n(x_n(z,s);q)(q;q)_\infty}{(-z)^n q^{n^2(1-s)}} = A_q\left(\frac{e^{2\lambda\pi i}}{z}\right) + r_{sw}(n|2), \tag{504}$$

*and*

$$|r_{sw}(n|2)| \le \frac{2A_q(-|z|^{-1})}{(q;q)_\infty} \left\{ q^{n/2} + \frac{q^{\lfloor n/2 \rfloor^2}}{|z|^{\lfloor n/2 \rfloor}} \right\} \tag{505}$$

*for $n$ is sufficiently large.*

3. *Assume that $\tau = 0$, if for any fixed real numbers $\beta$ and $\rho \ge 1$ there are infinitely many positive integers $n$ such that*

$$n\theta = m + \beta + b_n, \quad m \in \mathbb{Z}, \quad |b_n| < \frac{1}{n^\rho}, \tag{506}$$

*then,*

$$\frac{S_n(x_n(z,s);q)(q;q)_\infty}{(-z)^n q^{n^2(1-s)}} = A_q\left(\frac{e^{2\pi i\beta}}{z}\right) + e_{sw}(n|3), \tag{507}$$

*and*

$$|e_{sw}(n|3)| \le 24 \exp(q/((1-q)|z|)) \left\{ \frac{\log n}{n^\rho} + \frac{q^{n/2}}{(1-q)} + \frac{((q^{-1}-1)|z|)^{-k}}{j_n!} \right\} \tag{508}$$

*for $n$ is sufficiently large.*

(4) *Assume that* $-1 < \tau < 0$. *If for some fixed real numbers* $\lambda$ *and* $\lambda_1$ *there are infinite number of positive integers* $n$ *such that*

$$-n\tau = m+\lambda, \quad m\in\mathbb{N}, \quad n\theta = m_1+\lambda_1, \quad m_1\in\mathbb{Z}, \tag{509}$$

*then,*

$$S_n(x_n(z,s);q) = \frac{(-z)^n q^{n^2(1-s)+m(2\tau n+m)}}{(q;q)_\infty^2(-ze^{-2n\theta\pi i})^m}\left\{\theta_4\left(zq^{2\lambda}e^{-2\pi i\lambda_1};q\right)+r_{sw}(n|4)\right\}, \tag{510}$$

*and*

$$|r_{sw}(n|4)| \le 12\theta_3(|z|q^{2\lambda};q)\left\{\frac{q^{k_n}}{1-q}+|z|^{k_n}q^{k_n^2+2\lambda k_n}+\frac{q^{k_n^2-2\lambda k_n}}{|z|^{k_n}}\right\} \tag{511}$$

*for* $n$ *sufficiently large.*

(5) *Assume that* $-1 < \tau < 0$ , *if for any real numbers* $\beta$, $\lambda$ *and* $\rho \ge 1$ *there are infinitely many positive integers* $n$ *such that*

$$n\theta = m_1+\beta+b_n, \quad |b_n| < \frac{1}{n^\rho}, \quad m_1\in\mathbb{Z}, \tag{512}$$

*and*

$$-n\tau = m+\lambda, \quad m\in\mathbb{N}. \tag{513}$$

*Then,*

$$S_n(x_n(z,s);q) = \frac{(-z)^n q^{n^2(1-s)+m(2\tau n+m)}}{(q;q)_\infty^2(-ze^{-2n\theta\pi i})^m}\left\{\theta_4\left(zq^{2\lambda}e^{-2\beta\pi i};q\right)+r_{sw}(n|5)\right\}, \tag{514}$$

*and*

$$|r_{sw}(n|5)| \le 48\theta_3(|z|q^{2\lambda};q)\left\{\frac{q^{j_n^2-2\lambda j_n}}{|z|^{j_n}}+|z|^{j_n}q^{j_n^2+2\lambda j_n}+\frac{q^{k_n}}{1-q}+\frac{\log n}{n^\rho}\right\}, \tag{515}$$

*for* $n$ *sufficiently large.*

(6) *Assume that* $-1 < \tau < 0$, *if for any real numbers* $\beta$, $\lambda$ *and* $\rho \ge 1$ *there are infinitely many positive integers* $n$ *such that*

$$-n\tau = m+\beta+a_n, \quad |a_n| < \frac{1}{n^\rho}, \quad m\in\mathbb{N}, \tag{516}$$

*and*

$$n\theta = m_1+\lambda, \quad m_1\in\mathbb{Z}. \tag{517}$$

*Then,*

$$S_n(x_n(z,s);q) = \frac{(-z)^n q^{n^2(1-s)+m(2\tau n+m)}}{(q;q)_\infty^2(-ze^{-2n\theta\pi i})^m}\left\{\theta_4\left(zq^{2\beta}e^{-2\lambda\pi i};q\right)+r_{sw}(n|6)\right\}, \tag{518}$$

*and*

$$|r_{sw}(n|6)| \le 36\theta_3(|z|q^{2\beta};q)\left\{\frac{q^{j_n^2-2(\beta+1)j_n}}{|z|^{j_n}}+|z|^{j_n}q^{j_n^2+2(\beta-1)j_n}+\frac{q^{k_n}}{1-q}+\frac{\log n}{n^\rho}\right\}, \tag{519}$$

*for* $n$ *sufficiently large.*

(7) *Assume that* $-1 < \tau < 0$, *If for fixed real numbers* $\beta_1$, $\beta_2$ *and* $\rho > 0$ *there are infinitely many positive integers* $n$ *such that*

$$-\tau n = m + \beta_1 + a_n, \quad |a_n| < \frac{1}{n^\rho}, \quad m \in \mathbb{N}, \tag{520}$$

*and*

$$n\theta = m_1 + \beta_2 + b_n, \quad |b_n| < \frac{1}{n^\rho}, \quad m_1 \in \mathbb{Z}. \tag{521}$$

*Then,*

$$S_n(x_n(z,s);q) = \frac{(-z)^n q^{n^2(1-s)+m(2\tau n+m)}}{(q;q)_\infty^2(-ze^{-2n\theta\pi i})^m}\left\{\theta_4\left(zq^{2\beta_1}e^{-2\beta_2\pi i};q\right) + r_{sw}(n|7)\right\}, \tag{522}$$

*and*

$$|r_{sw}(n|7)| \le 156\theta_3(|z|q^{2\beta_1};q)\left\{\frac{q^{j_n^2-2(\beta_1+1)j_n}}{|z|^{j_n}} + |z|^{j_n}q^{j_n^2+2(\beta_1-1)j_n} + \frac{q^{k_n}}{1-q} + \frac{\log n}{n^\rho}\right\} \tag{523}$$

*for* $n$ *sufficiently large.*

In the following corollaries we assume that

$$z = e^{2\pi u}, \quad u \in \mathbb{R}. \tag{524}$$

If we let

$$q = \exp(-n^{-a}\pi), \quad 0 < a < \frac{1}{2}, \quad n \in \mathbb{N}, \tag{525}$$

then we have the following results for Stieltjes-Wigert polynomials:

**Corollary 6.2.** (1) *If*$\tau > 0$, *then*

$$\begin{aligned} &S_n(\exp 2\pi(u+(\tau+1)n^{1-a}); e^{-\pi n^{-a}}) \\ &\quad = \frac{\exp\left(2n\pi u + (2\tau+1)\pi n^{2-a} + \frac{\pi n^a}{6} - \frac{\pi n^{-a}}{24}\right)\left\{1+\mathcal{O}\left(e^{-4\pi n^a}\right)\right\}}{(-1)^n\sqrt{2n^a}}, \end{aligned} \tag{526}$$

*and*

$$\begin{aligned} &s_n(\exp 2\pi(u+(\tau+1)n^{1-a})\|e^{-\pi n^{-a}}) \\ &\quad = \frac{(-1)^n \exp\left(-\frac{\pi}{n^a}(n^a u + \tau n)^2 - \frac{\pi u}{2}\right)\left\{1+\mathcal{O}\left(e^{-4\pi n^a}\right)\right\}}{\sqrt{2\pi}\exp\left(\frac{\pi(\tau+2)n^{1-a}}{2} + \frac{\pi n^{-a}}{12} - \frac{\pi n^a}{12}\right)} \end{aligned} \tag{527}$$

*as* $n \to \infty$.

(2) *Assume that* $-1 < \tau < 0$ *and for some fixed real number* $\lambda$, *there are infinitely many positive integers* $n$ *such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}. \tag{528}$$

*Then,*

$$\begin{aligned} &S_n\left(\exp 2\pi(u+(\tau+1)n^{1-a}); e^{-\pi n^{-a}}\right) \\ &= \frac{\exp\left(\pi n^{-a}\left(n^a u+(\tau+1)n\right)^2\right)}{(-1)^{(1+\tau)n+\lambda} n^{a/2}\exp\left(\frac{\pi n^{-a}}{12}-\frac{\pi n^a}{12}\right)} \\ &\times\left\{\cos\pi(n^a u-\lambda)+\mathcal{O}\left(e^{-2\pi n^a}\right)\right\}, \end{aligned} \tag{529}$$

*and*

$$\begin{aligned} &s_n\left(\exp 2\pi(u+(\tau+1)n^{1-a}) \| e^{-\pi n^{-a}}\right) \\ &= \frac{(-1)^{(1+\tau)n+\lambda}\left\{\cos\pi(n^a u-\lambda)+\mathcal{O}\left(e^{-2\pi n^a}\right)\right\}}{\sqrt{\pi}\exp\left(\frac{\pi u}{2}+\frac{(\tau+2)\pi n^{1-a}}{2}+\frac{\pi n^{-a}}{8}\right)} \end{aligned} \tag{530}$$

*for $n$ sufficiently large.*

If we choose that

$$q=\exp\left(-\frac{\pi}{\gamma\log n}\right), \quad n\geq 2, \tag{531}$$

then we have the following:

**Corollary 6.3.** (1) *If $\tau>0$, then we have*

$$\begin{aligned} &S_n\left(\exp 2\pi\left(u+\frac{(\tau+1)n}{\gamma\log n}\right); \exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \\ &= \frac{\exp\left(2\pi nu+\frac{(2\tau+1)n^2}{\gamma\log n}+\frac{\gamma\pi\log n}{6}-\frac{\pi}{24\gamma\log n}\right)\left\{1+\mathcal{O}\left(n^{-4\pi\gamma}\right)\right\}}{(-1)^n\sqrt{2\gamma\log n}}, \end{aligned} \tag{532}$$

*and*

$$\begin{aligned} &s_n\left(\exp 2\pi\left(u+\frac{(\tau+1)n}{\gamma\log n}\right) \| \exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \\ &= \frac{\exp\left(-\frac{\pi}{\gamma\log n}(\gamma u\log n+\tau n)^2-\frac{\pi u}{2}\right)\left\{1+\mathcal{O}\left(n^{-4\pi\gamma}\right)\right\}}{(-1)^n\sqrt{2\pi}\exp\left(\frac{(\tau+2)n\pi}{2\gamma\log n}+\frac{\pi}{12\gamma\log n}-\frac{\gamma\pi\log n}{12}\right)} \end{aligned} \tag{533}$$

*for $n$ sufficiently large.*

(2) *Assume that $-1<\tau<0$ and for some fixed real number $\lambda$ there are infinitely many positive integers $n$ such that*

$$-n\tau=m+\lambda, \quad m\in\mathbb{N}. \tag{534}$$

*Then,*

$$\begin{aligned} &S_n\left(\exp 2\pi\left(u+\frac{(\tau+1)n}{\gamma\log n}\right); \exp\left(-\frac{\pi}{\gamma\log n}\right)\right) \\ &= \frac{(-1)^{(1+\tau)n+\lambda}n^{\gamma\pi/12}}{\sqrt{\gamma\log n}}\exp\left(\frac{\pi(u\gamma\log n+(\tau+1)n)^2}{\gamma\log n}-\frac{\pi}{12\gamma\log n}\right) \\ &\times\left\{\cos\pi(\gamma u\log n-\lambda)+\mathcal{O}\left(n^{-2\pi\gamma}\right)\right\}, \end{aligned} \tag{535}$$

*and*

$$s_n\left(\exp 2\pi\left(u+\frac{(\tau+1)n}{\gamma\log n}\right)\|\exp\left(-\frac{\pi}{\gamma\log n}\right)\right) = \frac{(-1)^{(1+\tau)n+\lambda}\left\{\cos\pi(\gamma u\log n-\lambda)+\mathcal{O}\left(n^{-2\pi\gamma}\right)\right\}}{\sqrt{\pi}\exp\left(\frac{\pi u}{2}+\frac{\pi(\tau+2)n}{2\gamma\log n}+\frac{\pi}{8\gamma\log n}\right)} \tag{536}$$

*for $n$ sufficiently large.*

(3) *Assume that $-1<\tau<0$, if for some fixed real numbers $\beta$, $\rho\ge 1$ and $\lambda$ there exist infinitely many positive integers $n$ such that*

$$-n\tau=m+\beta+a_n,\quad |a_n|<\frac{1}{n^\rho},\quad m\in\mathbb{N}, \tag{537}$$

*and let*

$$\gamma=\frac{4\rho}{9\pi}, \tag{538}$$

*then for each of such $n$, we have*

$$\begin{aligned} &S_n\left(\exp 2\pi\left(u+\frac{(\tau+1)n}{\gamma\log n}\right);\exp\left(-\frac{\pi}{\gamma\log n}\right)\right)\\ &=\frac{(-1)^{n-m}n^{\rho/27}}{\sqrt{\gamma\log n}}\exp\left(\frac{\pi(u\gamma\log n+(\tau+1)n)^2}{\gamma\log n}-\frac{\pi}{12\gamma\log n}\right)\\ &\times\left\{\cos\pi(\gamma u\log n-\beta)+\mathcal{O}\left(n^{-8\rho/9}\log n\right)\right\}, \end{aligned} \tag{539}$$

*and*

$$s_n\left(\exp 2\pi\left(u+\frac{(\tau+1)n}{\gamma\log n}\right)\|\exp\left(-\frac{\pi}{\gamma\log n}\right)\right) = \frac{(-1)^{n-m}\left\{\cos\pi(\gamma u\log n-\beta)+\mathcal{O}\left(n^{-8\rho/9}\log n\right)\right\}}{\sqrt{\pi}\exp\left(\frac{\pi u}{2}+\frac{\pi(\tau+2)n}{2\gamma\log n}+\frac{\pi}{8\gamma\log n}\right)} \tag{540}$$

*for $n$ sufficiently large.*

6.2. **Proofs for Theorem 6.1.** In the first three proofs we will use the inequalities

$$0<\frac{(q;q)_\infty}{(q;q)_{n-k}}\le 1 \tag{541}$$

for $0\le k\le n$ and

$$\left|\frac{(q;q)_\infty}{(q;q)_{n-k}}-1\right|\le\frac{2q^{1+n/2}}{1-q} \tag{542}$$

for $0\le k\le\left\lfloor\frac{n}{2}\right\rfloor-1$ for $n$ sufficiently large, which is a directly from Lemma 1.1.

In the last four proofs we have

$$-\tau n=m+c_n,\quad m\in\mathbb{N},\quad n\theta=m_1+d_n,\quad m_1\in\mathbb{Z}. \tag{543}$$

Then,

$$\begin{aligned}(544)\qquad \frac{S_n(x_n(z,s);q)}{(-z)^n q^{n^2(1-s)}} &= \sum_{k=0}^{n} \frac{q^{k^2} e^{2nk\theta\pi i}}{(q;q)_k (q;q)_{n-k}} \left(-\frac{q^{2\tau n}}{z}\right)^k \\ &= \sum_{k=0}^{m} \frac{q^{k^2} e^{2nk\theta\pi i}}{(q;q)_k (q;q)_{n-k}} \left(-\frac{q^{2\tau n}}{z}\right)^k + \sum_{k=m+1}^{n} \frac{q^{k^2} e^{2nk\theta\pi i}}{(q;q)_k (q;q)_{n-k}} \left(-\frac{q^{2\tau n}}{z}\right)^k \\ &= s_1 + s_2.\end{aligned}$$

We reverse the summation order in $s_1$ to obtain

$$(545)\qquad \frac{s_1 (q;q)_\infty^2 (-ze^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}} = \sum_{k=0}^{m} q^{k^2} \left(-zq^{2c_n} e^{-2\pi i d_n}\right)^k e(k,n),$$

and

$$(546)\qquad e(k,n) = \frac{(q;q)_\infty^2}{(q;q)_{m-k}(q;q)_{n-m+k}}.$$

It is clear that

$$(547)\qquad |e(k,n)| \le 1$$

for $0 \le k \le m$. Using Lemma 1.1 to expand and estimate each term of

$$(548)\qquad e(k,n) \quad -1 = \quad \{r_1(q; m-k) + 1\}\{r_1(q; n-m+k) + 1\} - 1,$$

to get

$$(549)\qquad |e(k,n) - 1| \le \frac{6q^{k_n+2}}{1-q}$$

for $0 \le k \le k_n - 1$ and $n$ sufficiently large.

In sum $s_2$ we shift summation index from $k$ to $k+m$ to get

$$(550)\qquad \frac{s_2 (q;q)_\infty^2 (-ze^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}} = \sum_{k=1}^{n-m} q^{k^2} \left(-z^{-1} q^{-2c_n} e^{2\pi i d_n}\right)^k f(k,n),$$

and

$$(551)\qquad f(k,n) = \frac{(q;q)_\infty^2}{(q;q)_{m+k}(q;q)_{n-m-k}}$$

for $1 \le k \le n-m$. Hence

$$(552)\qquad |f(k,n)| \le 1$$

for $1 \le k \le n-m$. Expand and apply Lemma 1.1 to each terms of

$$(553)\qquad f(k,n) \quad -1 = \quad \{r_1(q; m+k) + 1\}\{r_1(q; n-m-k) + 1\} - 1$$

to obtain

$$(554)\qquad |f(k,n) - 1| \le \frac{6q^{k_n+2}}{1-q}$$

for $1 \le k \le k_n - 1$ and $n$ sufficiently large.

The rest are similar to the corresponding proofs for the Ismail-Masson polynomials.

6.3. **Proof for Corollary 6.2.** In this case we have

$$x_n(z,s) = \exp 2\pi(u+(\tau+1)n^{1-a}) \tag{555}$$

and

$$w_{sw}(x_n(z,s)) = \frac{n^{a/2}}{\pi\sqrt{2}} \exp\left(-2\pi n^{-a}(n^a u + (\tau+1)n + \frac{1}{4})^2\right). \tag{556}$$

Clearly,

$$(q;q)_\infty = \sqrt{2n^a}\exp\left(\frac{\pi n^{-a}}{24} - \frac{\pi n^a}{6}\right)\left\{1+\mathcal{O}\left(e^{-4\pi n^a}\right)\right\}, \tag{557}$$

$$\frac{1}{(q;q)_\infty} = \frac{\exp\left(\frac{\pi n^a}{6} - \frac{\pi n^{-a}}{24}\right)\left\{1+\mathcal{O}\left(e^{-4\pi n^a}\right)\right\}}{\sqrt{2n^a}}, \tag{558}$$

$$q^n(q;q)_n = \sqrt{2n^a}\exp\left(-\pi n^{1-a} + \frac{\pi n^{-a}}{24} - \frac{\pi n^a}{6}\right)\left\{1+\mathcal{O}\left(e^{-4\pi n^a}\right)\right\}, \tag{559}$$

then,

$$\begin{aligned} &\sqrt{q^n(q;q)_n w_{sw}(x_n(z,s))} \\ &= \sqrt{\frac{n^a}{\pi}}\exp\left(\frac{\pi n^{-a}}{48} - \frac{\pi n^a}{12} - \frac{\pi n^{1-a}}{2}\right) \\ &\times \exp\left(-\pi n^{-a}(n^a u + (\tau+1)n + \frac{1}{4})^2\right)\left\{1+\mathcal{O}\left(e^{-4\pi n^a}\right)\right\} \end{aligned} \tag{560}$$

as $n\to\infty$.

Notice that

$$\begin{aligned} \theta_4\left(zq^{2\lambda};q\right) &= \theta_4\left(ui - i\lambda n^{-a} \mid n^{-a}i\right) \\ &= n^{a/2}\exp\left(\frac{\pi(n^a u-\lambda)^2}{n^a}\right)\theta_2\left(n^a u - \lambda \mid n^a i\right) \\ &= 2n^{a/2}\exp\left(\frac{\pi(n^a u-\lambda)^2}{n^a} - \frac{\pi n^a}{4}\right) \\ &\times \cos\pi(n^a u - \lambda)\left\{1+\mathcal{O}\left(e^{-2\pi n^a}\right)\right\}, \end{aligned} \tag{561}$$

and

$$\begin{aligned} \theta_3\left(zq^{2\lambda};q\right) &= \theta_3\left(ui - i\lambda n^{-a} \mid n^{-a}i\right) \\ &= n^{a/2}\exp\left(\frac{\pi(n^a u-\lambda)^2}{n^a}\right)\theta_3\left(n^a u - \lambda \mid n^a i\right) \\ &= 2n^{a/2}\exp\left(\frac{\pi(n^a u-\lambda)^2}{n^a}\right)\left\{1+\mathcal{O}\left(e^{-\pi n^a}\right)\right\} \end{aligned} \tag{562}$$

as $n\to\infty$. The proof for case 1 follows from (501) and (502), while the proof for case 2 comes from (510) and (511).

6.4. **Proof for Corollary 6.3.** In this case we have

$$x_n(z,s)=\exp 2\pi\left(u+\frac{(\tau+1)n}{\gamma\log n}\right), \tag{563}$$

and

$$w_{sw}(x_n(z,s))=\frac{1}{\pi}\sqrt{\frac{\gamma\log n}{2}}\exp\left(-\frac{2\pi}{\gamma\log n}(\gamma u\log n+(\tau+1)n+\frac{1}{4})^2\right). \tag{564}$$

It is clear that

$$\begin{aligned}(q;q)_\infty &= \sqrt{2\gamma\log n}\exp\left(\frac{\pi}{24\gamma\log n}-\frac{\gamma\pi\log n}{6}\right)\left\{1+\mathcal{O}\left(n^{-4\pi\gamma}\right)\right\}, && (565)\\ \frac{1}{(q;q)_\infty} &= \frac{1}{\sqrt{2\gamma\log n}}\exp\left(\frac{\gamma\pi\log n}{6}-\frac{\pi}{24\gamma\log n}\right)\left\{1+\mathcal{O}\left(n^{-4\pi\gamma}\right)\right\} && (566)\end{aligned}$$

as $n\to\infty$.

$$\begin{aligned}&\sqrt{q^n(q;q)_n w_{sw}(x_n(z,s))}\\ &=\sqrt{\frac{\gamma\log n}{\pi}}\exp\left(\frac{\pi}{48\gamma\log n}-\frac{\pi n}{2\gamma\log n}-\frac{\gamma\pi\log n}{12}\right)\\ &\times\exp\left(-\frac{\pi}{\gamma\log n}(\gamma u\log n+(\tau+1)n+\frac{1}{4})^2\right)\left\{1+\mathcal{O}\left(n^{-4\pi\gamma}\right)\right\}\end{aligned} \tag{567}$$

as $n\to\infty$.

(568)

$$\begin{aligned}\theta_4(zq^{2\lambda};q)&=\theta_4\left(ui-\frac{\lambda i}{\gamma\log n}\mid\frac{i}{\gamma\log n}\right)\\ &=\sqrt{\gamma\log n}\exp\frac{\pi(\gamma u\log n-\lambda)^2}{\gamma\log n}\theta_2\left(\gamma u\log n-\lambda\mid i\gamma\log n\right)\\ &=\sqrt[24]{\frac{(\gamma\log n)^2}{n^{\pi\gamma}}}\exp\frac{\pi(\gamma u\log n-\lambda)^2}{\gamma\log n}\cos\pi(\gamma u\log n-\lambda)\left\{1+\mathcal{O}\left(n^{-2\pi\gamma}\right)\right\},\end{aligned}$$

and

$$\begin{aligned}\theta_3(zq^{2\lambda};q)&=\theta_3\left(ui-\frac{\lambda i}{\gamma\log n}\mid\frac{i}{\gamma\log n}\right)\\ &=\sqrt{\gamma\log n}\exp\frac{\pi(\gamma u\log n-\lambda)^2}{\gamma\log n}\theta_3\left(\gamma u\log n-\lambda\mid i\gamma\log n\right)\\ &=\sqrt{\gamma\log n}\exp\frac{\pi(\gamma u\log n-\lambda)^2}{\gamma\log n}\left\{1+\mathcal{O}\left(n^{-\pi\gamma}\right)\right\},\end{aligned} \tag{569}$$

as $n\to\infty$. The proofs for three cases follow from the formulas (501), (510) and (518) respectively.

## 7. $q$-LAGUERRE ORTHOGONAL POLYNOMIALS

The $q$-Laguerre orthogonal polynomials $\left\{L_n^{(\alpha)}(x;q)\right\}_{n=0}^{\infty}$ are defined as [13]

$$L_n^{(\alpha)}(x;q)=\sum_{k=0}^{n}\frac{q^{k^2+\alpha k}(-x)^k(q^{\alpha+1};q)_n}{(q;q)_k(q,q^{\alpha+1};q)_{n-k}}, \tag{570}$$

for $\alpha > -1$. For any nonzero complex number $z$, let

$$s = 2\tau + 2 + i\frac{2\theta\pi}{\log q}, \quad x_n(z,s,\alpha) = zq^{-ns-\alpha}, \quad \tau, \theta \in \mathbb{R}, \tag{571}$$

reverse the summation order to get

$$\frac{L_n^{(\alpha)}(x_n(z,s,\alpha);q)}{(-z)^n q^{n^2(1-s)}} = \sum_{k=0}^{n} \frac{q^{k^2} e^{2nk\theta\pi i}}{(q;q)_k (q;q)_{n-k}} \frac{(q^{\alpha+1};q)_n}{(q^{\alpha+1};q)_{n-k}} \left(-\frac{q^{2\tau n}}{z}\right)^k, \tag{572}$$

then

$$\left|L_n^{(\alpha)}(x_n(z,s,\alpha);q)\right| \le \frac{|z|^n \sum_{k=0}^{\infty} \frac{q^{k^2}\left(|z|^{-1}q^{2\tau n}\right)^k}{(q;q)_k}}{(q;q)_\infty q^{n^2(2\tau+1)}} \le \frac{|z|^n A_q\left(-|z|^{-1}q^{2\tau n}\right)}{(q;q)_\infty q^{n^2(2\tau+1)}}, \tag{573}$$

or

$$\left|L_n^{(\alpha)}(x_n(z,s,\alpha);q)\right| \le \frac{(-\sqrt{q};q)_\infty}{(q;q)_\infty} \frac{|z|^{(2\tau+1)n} \exp\left(-\frac{\log^2|z|}{2\log q}\right)}{q^{(2\tau^2+2\tau+1)n^2}}. \tag{574}$$

### 7.1. **Asymptotic Formulas $q$-Laguerre Polynomials.**

**Theorem 7.1.** *Assume that $\alpha > -1$, given any complex numbers $z \neq 0$, let $s$ and $x_n(z,s,\alpha)$ be defined as in (601) and let*

$$j_n = \left\lfloor \frac{q^4 \log n}{\log q^{-1}} \right\rfloor, \quad k_n = \min\left\{ \left\lfloor \frac{(1+\tau)n}{2} \right\rfloor, \left\lfloor \frac{-\tau n}{2} \right\rfloor \right\}, \tag{575}$$

*we have the following results for $q$-Laguerre polynomials:*

(1) *Assume that $\tau > 0$, we have*

$$\frac{L_n^{(\alpha)}(x_n(z,s,\alpha);q)(q;q)_\infty}{(-z)^n q^{n^2(1-s)}} = 1 + r_{ql}(n|1) \tag{576}$$

*with*

$$|r_{ql}(n|1)| \le \frac{q^{2\tau n+1} \exp\left(q^{2\tau n+1}/((1-q)|z|)\right)}{(1-q)|z|}. \tag{577}$$

(2) *Assume that $\tau = 0$, if for some fixed real number $\lambda$ there are infinitely positive integers $n$ such that*

$$n\theta = m + \lambda, \quad m \in \mathbb{Z}, \tag{578}$$

*then,*

$$\frac{L_n^{(\alpha)}(x_n(z,s,\alpha);q)(q;q)_\infty}{(-z)^n q^{n^2(1-s)}} = A_q\left(\frac{e^{2\lambda\pi i}}{z}\right) + r_{ql}(n|2), \tag{579}$$

*and*

$$|r_{ql}(n|2)| \le \frac{14A_q(-|z|^{-1})}{(q;q)_\infty} \left\{ q^{n/2} + \frac{q^{\lfloor n/2 \rfloor^2}}{|z|^{\lfloor n/2 \rfloor}} \right\} \tag{580}$$

*for $n$ sufficiently large.*

(3) *Assume that $\tau = 0$, if for some fixed real numbers $\beta$ and $\rho \ge 1$ there are infinitely many positive integers $n$ such that*

$$n\theta = m + \beta + b_n \quad |b_n| < \frac{1}{n^{\rho}}, \quad m \in \mathbb{Z}, \tag{581}$$

*then,*

$$\frac{L_n^{(\alpha)}(x_n(z,s,\alpha);q)(q;q)_\infty}{(-z)^n q^{n^2(1-s)}} = A_q\left(\frac{e^{2\pi i\beta}}{z}\right) + e_{ql}(n|3), \tag{582}$$

*and*

$$|e_{ql}(n|3)| \le 24\exp\left(\frac{q}{(1-q)z}\right)\left\{\frac{\log n}{n^{\rho}} + \frac{q^{n/2}}{1-q} + \frac{\left((q^{-1}-1)|z|\right)^{-j_n}}{j_n!}\right\} \tag{583}$$

*for $n$ sufficiently large.*

(4) *Assume that $-1 < \tau < 0$, if for some fixed real numbers $\lambda$ and $\lambda_1$ there are infinite number of positive integers $n$ such that*

$$-n\tau = m + \lambda, \quad m \in \mathbb{N}, \quad n\theta = m_1 + \lambda_1, \quad m_1 \in \mathbb{Z}, \tag{584}$$

*then,*

$$L_n^{(\alpha)}(x_n(z,s,\alpha);q) = \frac{(-z)^n q^{n^2(1-s)+m(2\tau n+m)}}{(q;q)_\infty^2\left(-ze^{-2n\theta\pi i}\right)^m}\left\{\theta_4\left(zq^{2\lambda}e^{-2\pi i\lambda_1};q\right) + r_{ql}(n|4)\right\}, \tag{585}$$

*and*

$$|r_{ql}(n|4)| \le 60\theta_3\left(|z|q^{2\lambda};q\right)\left\{\frac{q^{k_n}}{1-q} + |z|^{-k_n}q^{k_n^2-2\lambda k_n} + |z|^{k_n}q^{k_n^2+2\lambda k_n}\right\} \tag{586}$$

*for $n$ sufficiently large.*

(5) *Assume that $-1 < \tau < 0$ , if for some fixed real numbers $\beta$, $\lambda$ and $\rho \ge 1$ there are infinitely many positive integers $n$ such that*

$$n\theta = m_1 + \beta + b_n, \quad |b_n| < \frac{1}{n^{\rho}}, \quad m_1 \in \mathbb{Z}, \quad -n\tau = m + \lambda, \quad m \in \mathbb{N}, \tag{587}$$

*then,*

$$L_n^{(\alpha)}(x_n(z,s,\alpha);q) = \frac{(-z)^n q^{n^2(1-s)+m(2\tau n+m)}}{(q;q)_\infty^2(-ze^{-2n\theta\pi i})^m}\left\{\theta_4\left(zq^{2\lambda}e^{-2\beta\pi i};q\right) + r_{ql}(n|5)\right\}, \tag{588}$$

*and*

$$|r_{ql}(n|5)| \le 60\theta_3(|z|q^{2\lambda};q)\left\{|z|^{j_n}q^{j_n^2+2\lambda j_n} + |z|^{-j_n}q^{j_n^2-2\lambda j_n} + \frac{q^{k_n}}{1-q} + \frac{\log n}{n^{\rho}}\right\} \tag{589}$$

*for $n$ sufficiently large.*

(6) *Assume that $-1 < \tau < 0$, if for any fixed real numbers $\beta$,$\lambda$ and $\rho \ge 1$ there are infinitely many positive integers $n$ such that*

$$-n\tau = m + \beta + a_n, \quad |a_n| < \frac{1}{n^{\rho}}, \quad m \in \mathbb{N}, \quad n\theta = m_1 + \lambda, \quad m_1 \in \mathbb{Z}, \tag{590}$$

*then,*

(591)
$$L_n^{(\alpha)}(x_n(z,s,\alpha);q)=\frac{(-z)^n q^{n^2(1-s)+m(2\tau n+m)}}{(q;q)_\infty^2(-ze^{-2n\theta\pi i})^m}\left\{\theta_4\left(zq^{2\beta}e^{-2\lambda\pi i};q\right)+r_{ql}(n|6)\right\},$$

*and*

(592)
$$|r_{ql}(n|6)|\le 180\theta_3(|z|q^{2\beta};q)\left\{|z|^{-j_n}q^{j_n^2-2(\beta+1)j_n}+|z|^{j_n}q^{j_n^2+2(\beta-1)j_n}+\frac{q^{k_n}}{1-q}+\frac{\log n}{n^\rho}\right\}$$

*for $n$ sufficiently large.*

(7) *Assume that $-1<\tau<0$, if for some fixed real numbers $\beta_1$, $\beta_2$ and $\rho>0$ there are infinitely many positive integers $n$ such that*

$$-\tau n=m+\beta_1+a_n,\quad |a_n|<\frac{1}{n^\rho},\quad m\in\mathbb{N}, \tag{593}$$

*and*

$$n\theta=m_1+\beta_2+b_n,\quad |b_n|<\frac{1}{n^\rho},\quad m_1\in\mathbb{Z}, \tag{594}$$

*then,*

(595)
$$L_n^{(\alpha)}(x_n(z,s,\alpha);q)=\frac{(-z)^n q^{n^2(1-s)+m(2\tau n+m)}}{(q;q)_\infty^2(-ze^{-2n\theta\pi i})^m}\left\{\theta_4\left(zq^{2\beta_1}e^{-2\beta_2\pi i};q\right)+r_{ql}(n|7)\right\},$$

*and*

$$|r_{ql}(n|7)|\le 180\left\{|z|^{-j_n}q^{j_n^2-2(\beta_1+1)j_n}+|z|^{j_n}q^{j_n^2+2(\beta_1-1)j_n}+\frac{q^{k_n}}{1-q}+\frac{\log n}{n^\rho}\right\} \tag{596}$$

*for $n$ sufficiently large.*

In the following corollaries we assume that

$$z=e^{2\pi u},\quad u\in\mathbb{R}. \tag{597}$$

If we let

$$q=\exp(-n^{-a}\pi),\quad 0<a<\frac{1}{2},\quad n\in\mathbb{N}, \tag{598}$$

then we have the following results for $q$-Laguerre orthogonal polynomials:

**Corollary 7.2.** (1) *If$\tau>0$, then*

$$\begin{aligned}&L_n^{(\alpha)}\left(\exp 2\pi(u+(\tau+1)n^{1-a}+\alpha n^{-a}/2);\exp(-n^{-a}\pi)\right)\\&=\frac{\exp 2n\pi\left(u+(\tau+\frac{1}{2})n^{1-a}\right)\left\{1+\mathcal{O}(e^{-4\pi n^a})\right\}}{(-1)^n\sqrt{2n^a}\exp\left(-\frac{\pi}{24}(n^{-a}-4n^a)\right)}\end{aligned} \tag{599}$$

*as $n\to\infty$.*

(2) *Assume that $-1<\tau<0$ and for some fixed real number $\lambda$ there are infinitely many positive integers $n$ such that*

$$-n\tau=m+\lambda,\quad m\in\mathbb{N}. \tag{600}$$

*Then,*

$$L_n^{(\alpha)}(\exp 2\pi(u+(\tau+1)n^{1-a}+\alpha n^{-a}/2);\exp(-n^{-a}\pi)) = \frac{\exp\left(\pi n^{-a}(n^a u+(\tau+1)n)^2\right)\left\{\cos\pi(n^a u-\lambda)+\mathcal{O}\left(e^{-2\pi n^a}\right)\right\}}{(-1)^{(1+\tau)n+\lambda}\sqrt{n^a}\exp\left(\frac{\pi n^{-a}}{24}+\frac{\pi n^a}{12}\right)} \tag{601}$$

*for $n$ sufficiently large.*

If we choose

$$q=\exp\left(-\frac{\pi}{\gamma\log n}\right),\quad n\geq 2, \tag{602}$$

then we have the following:

**Corollary 7.3.** (1) *If $\tau>0$, then we have*

$$\begin{aligned}&L_n^{(\alpha)}\left(\exp\frac{2\pi(\gamma u\log n+(\tau+1)n+\alpha/2)}{\gamma\log n};\exp\left(-\frac{\pi}{\gamma\log n}\right)\right)\\&=\frac{n^{\pi\gamma/6}\exp\left(2n\pi u+\frac{(2\tau+1)n^2\pi}{\gamma\log n}-\frac{\pi}{24\gamma\log n}\right)}{(-1)^n\sqrt{2\gamma\log n}}\left\{1+\mathcal{O}\left(n^{-4\pi\gamma}\right)\right\}\end{aligned} \tag{603}$$

*for $n$ sufficiently large.*

(2) *Assume that $-1<\tau<0$ and for some fixed real number $\lambda$ there are infinitely many positive integers $n$ such that*

$$-n\tau=m+\lambda,\quad m\in\mathbb{N}, \tag{604}$$

*then,*

$$\begin{aligned}&L_n^{(\alpha)}\left(\exp\left(\frac{2\pi(\gamma u\log n+(\tau+1)n+\alpha/2)}{\gamma\log n}\right);\exp\left(-\frac{\pi}{\gamma\log n}\right)\right)\\&=\frac{n^{\pi\gamma/12}\exp\left(\frac{\pi(u\gamma\log n+(\tau+1)n)^2}{\gamma\log n}-\frac{\pi}{12\gamma\log n}\right)\left\{\cos\pi(\gamma u\log n-\lambda)+\mathcal{O}\left(n^{-2\pi\gamma}\right)\right\}}{(-1)^{(1+\tau)n+\lambda}\sqrt{\gamma\log n}}\end{aligned} \tag{605}$$

*for $n$ sufficiently large.*

(3) *Assume that $-1<\tau<0$, if for some fixed real numbers $\beta$, $\rho\geq 1$ and $\lambda$ there exist infinitely many positive integers $n$ such that*

$$-n\tau=m+\beta+a_n,\quad |a_n|<\frac{1}{n^\rho},\quad m\in\mathbb{N}, \tag{606}$$

*and let*

$$\gamma=\frac{4\rho}{9\pi}, \tag{607}$$

*then for each of such $n$, we have*

$$\begin{aligned}&L_n^{(\alpha)}\left(\exp\left(\frac{2\pi(\gamma u\log n+(\tau+1)n+\alpha/2)}{\gamma\log n}\right);\exp\left(-\frac{\pi}{\gamma\log n}\right)\right)\\&=\frac{\exp\left(\frac{\pi(u\gamma\log n+(\tau+1)n)^2}{\gamma\log n}-\frac{\pi}{12\gamma\log n}\right)}{(-1)^{n-m}n^{-\rho/27}\sqrt{\gamma\log n}}\left\{\cos\pi(\gamma u\log n-\beta)+\mathcal{O}\left(\frac{\log n}{n^{8\rho/9}}\right)\right\}\end{aligned} \tag{608}$$

*for $n$ sufficiently large.*

7.2. **Proofs for $q$-Laguerre Polynomial Case.** In the first three proofs we will use the inequalities

$$0 < \frac{(q;q)_\infty}{(q;q)_{n-k}} \frac{(q^{\alpha+1};q)_n}{(q^{\alpha+1};q)_{n-k}} \le 1 \tag{609}$$

for $0 \le k \le n$ and

$$\left| \frac{(q;q)_\infty}{(q;q)_{n-k}} \frac{(q^{\alpha+1};q)_n}{(q^{\alpha+1};q)_{n-k}} - 1 \right| \le \frac{14 q^{n/2}}{1-q} \tag{610}$$

for $0 \le k \le \lfloor \frac{n}{2} \rfloor$-1 and $n$ sufficiently large. This can be seen by expanding

(611)

$$\frac{(q;q)_\infty}{(q;q)_{n-k}} \frac{(q^{\alpha+1};q)_n}{(q^{\alpha+1};q)_{n-k}} - 1 = \left\{ r_1(q;n-k) + 1 \right\} \left\{ r_2(q^{\alpha+1};n) + 1 \right\} \left\{ r_1(q^{\alpha+1};n-k) + 1 \right\} - 1$$

and estimating each term by Lemma 1.1.

Assume that

$$-\tau n = m + c_n, \quad m \in \mathbb{N}, \quad n\theta = m_1 + d_n, \quad m_1 \in \mathbb{Z}, \tag{612}$$

then,

$$\begin{aligned} \frac{L_n^{(\alpha)}(x_n(z,s,\alpha);q)}{(-z)^n q^{n^2(1-s)}} &= \sum_{k=0}^{n} \frac{q^{k^2} e^{2nk\theta\pi i}}{(q;q)_k (q;q)_{n-k}} \frac{(q^{\alpha+1};q)_n}{(q^{\alpha+1};q)_{n-k}} \left( -\frac{q^{2\tau n}}{z} \right)^k \\ &= \sum_{k=0}^{m} \frac{q^{k^2} e^{2nk\theta\pi i}}{(q;q)_k (q;q)_{n-k}} \frac{(q^{\alpha+1};q)_n}{(q^{\alpha+1};q)_{n-k}} \left( -\frac{q^{2\tau n}}{z} \right)^k \\ &+ \sum_{k=m+1}^{n} \frac{q^{k^2} e^{2nk\theta\pi i}}{(q;q)_k (q;q)_{n-k}} \frac{(q^{\alpha+1};q)_n}{(q^{\alpha+1};q)_{n-k}} \left( -\frac{q^{2\tau n}}{z} \right)^k = s_1 + s_2. \end{aligned} \tag{613}$$

We reverse the summation order in $s_1$ to obtain

$$\frac{s_1 (q;q)_\infty^2 (-z e^{-2n\theta\pi i})^m}{q^{m(2\tau n + m)}} = \sum_{k=0}^{m} q^{k^2} \left( -z q^{2c_n} e^{-2\pi i d_n} \right)^k e(k,n), \tag{614}$$

and

$$e(k,n) = \frac{(q;q)_\infty^2 (q^{\alpha+1};q)_n}{(q;q)_{m-k} (q;q)_{n-m+k} (q^{\alpha+1};q)_{n-m+k}}. \tag{615}$$

Clearly,

$$|e(k,n)| \le 1 \tag{616}$$

for $0 \le k \le m$. Expand and estimate each term of

(617)

$$e(k,n) - 1 = \left\{ r_2\left(q^{\alpha+1};n\right) + 1 \right\} \left\{ r_1\left(q;m-k\right) + 1 \right\} \left\{ r_1\left(q;n-m+k\right) + 1 \right\} \left\{ r_1\left(q^{\alpha+1};n-m+k\right) + 1 \right\} - 1$$

by Lemma 1.1 to get

$$|e(k,n) - 1| \le \frac{30 q^{k_n+1}}{1-q} \tag{618}$$

for $0 \le k \le k_n - 1$.

In sum $s_2$ we shift summation from $k$ to $k+m$ to obtain

$$\frac{s_2(q;q)_\infty^2(-ze^{-2n\theta\pi i})^m}{q^{m(2\tau n+m)}} = \sum_{k=1}^{n-m} q^{k^2}\left(-z^{-1}q^{-2c_n}e^{2\pi i d_n}\right)^k f(k,n), \tag{619}$$

and

$$f(k,n) = \frac{(q;q)_\infty^2(q^{\alpha+1};q)_n}{(q;q)_{m+k}(q;q)_{n-m-k}(q^{\alpha+1};q)_{n-m-k}} \tag{620}$$

for $1 \le k \le n-m$. Hence

$$|f(k,n)| \le 1 \tag{621}$$

for $1 \le k \le n-m$. Apply Lemma 1.1 to each term of

$$f(k,n)-1 = \left\{r_2\left(q^{\alpha+1};n\right)+1\right\}\left\{r_1\left(q;m+k\right)+1\right\}\left\{r_1\left(q;n-m-k\right)+1\right\}\left\{r_1\left(q^{\alpha+1};n-m-k\right)+1\right\}-1, \tag{622}$$

we obtain

$$|f(k,n)-1| \le \frac{30q^{k_n+1}}{1-q} \tag{623}$$

for $1 \le k \le k_n - 1$.

The rest of the proofs are very similar to the corresponding proofs for Ismail-Masson polynomials and we will not repeat them here.

## REFERENCES

[1] N. I. Akhiezer, The Classical Moment Problem and Some Related Questions in Analysis, English translation, Oliver and Boyed, Edinburgh, 1965.

[2] Tom Apostol, Modular Functions and Dirichlet Series in Number Theory, Second Edition, Springer-Verlag, New York, 1990.

[3] G. E. Andrews, q-series: Their development and application in analysis, number theory, combinatorics, physics, and computer algebra, CBMS Regional Conference Series, number 66, American Mathematical Society, Providence, R.I. 1986.

[4] G. E. Andrews, Ramanujan's "Lost" Note book VIII: The entire Rogers-Ramnujan function, Advances in Math. 191 (2005), 393–407.

[5] G. E. Andrews, Ramanujan's "Lost" Note book IX: The entire Rogers-Ramnujan function, Advances in Math. 191 (2005), 408–422.

[6] G. E. Andrews, R. A. Askey, and R. Roy, Special Functions, Cambridge University Press, Cambridge, 1999.

[7] P. Deift, Orthogonal Polynomials and Random Matrices: a Riemann-Hilbert Approach, American Mathematical Society, Providence, 2000.

[8] P. Deift, T. Kriecherbauer, K. T-R. McLaughlin, S. Venakides, and X. Zhou, Strong asymptotics of orthogonal polynomials with respect to exponential weights, Comm. Pure Appl. Math. 52 (1999), 1491–1552.

[9] G. Gasper and M. Rahman, Basic Hypergeometric Series, second edition Cambridge University Press, Cambridge, 2004.

[10] W. K. Hayman, On the zeros of a q-Bessel function, Contemporary Mathematics, volume 382, American Mathematical Society, Providence, 2005, 205–216.

[11] Hua Loo Keng, Introduction to Number Theory, Springer-Verlag, Berlin Heidelberg New York, 1982.

[12] M. E. H. Ismail, Asymptotics of q-orthogonal polynomials and a q-Airy function, Internat. Math. Res. Notices 2005 No 18 (2005), 1063–1088.

[13] M. E. H. Ismail, Classical and Quantum Orthogonal Polynomials in one Variable, Cambridge University Press, Cambridge, 2005.

[14] M. E. H. Ismail and X. Li, Bounds for extreme zeros of orthogonal polynomials, Proc. Amer. Math. Soc. 115 (1992), 131–140.

[15] M. E. H. Ismail and D. R. Masson, q-Hermite polynomials, biorthogonal rational functions, Trans. Amer. Math. Soc. 346 (1994), 63–116.
[16] M. E. H. Ismail and C. Zhang, Zeros of entire functions and a problem of Ramanujan, Advances in Math., (2007), to appear.
[17] M. E. H. Ismail and R. Zhang, Scaled asymptotics for q-polynomials, Comptes Rendus, submitted.
[18] M. E. H. Ismail and R. Zhang, Chaotic and Periodic Asymptotics for q-Orthogonal Polynomials, joint with Mourad E.H. Ismail, International Mathematics Research Notices, accepted.
[19] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta, Y. Yamada, Hypergeometric solutions to the q-Painlev\'{e} equations, Internat. Math. Res. Notices 47 (2004), 2497–2521.
[20] R. Koekoek and R. Swarttouw, The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogues, Reports of the Faculty of Technical Mathematics and Informatics no. 98-17, Delft University of Technology, Delft, 1998.
[21] M. L. Mehta, Random Matrices, third edition, Elsevier, Amsterdam, 2004.
[22] S. Ramanujan, The Lost Notebook and Other Unpublished Papers (Introduction by G. E. Andrews), Narosa, New Delhi, 1988.
[23] Hans Rademarcher, *Topics in Analytic Number Theory*, Die Grundlehren der mathematischen Wissenschaften, Bd. 169, Springer-Verlag, New York-Heidelberg-Berlin, 1973. Z. 253. 10002.
[24] E. B. Saff and V. Totik, Logarithmic Potentials With External Fields, Springer-Verlag, New York, 1997.
[25] G. Szeg\H{o}, Orthogonal Polynomials, Fourth Edition, Amer. Math. Soc., Providence, 1975.
[26] R. Wong, Asymptotic Approximations of Intehrals, Academic Press, Boston, 1989.
[27] E. T. Whittaker and G. N. Watson, A Course of Modern Analysis, fourth edition, Cambridge University Press, Cambridge, 1927. Hua Loo Keng, Introduction to Number Theory, Springer-Verlag, Berlin Heidelberg New York, 1982.
[28] Wikipedia.org, *http://en.wikipedia.org/wiki/Liouville_ number.*

*Current address*: School of Mathematics, Guangxi Normal University, Guilin City, Guangxi 541004, P. R. China.

*E-mail address*: `ruimingzhang@yahoo.com`